

3.4 Stock summaries

3.4.1 Cod in Subareas I and II (Northeast Arctic cod)

State of the stock

Spawning biomass in relation to precautionary limits	Fishing mortality in relation to precautionary limits/management plan	Fishing mortality in relation to highest yield	Fishing mortality in relation to agreed target	Comment
Full reproductive capacity	Harvested sustainably	Appropriate	Appropriate	

Based on the most recent estimates of SSB and fishing mortality, ICES classifies the stock as having full reproductive capacity and being harvested sustainably. The SSB has been above B_{pa} since 2002. Fishing mortality was reduced from well above F_{lim} in 1999 to below F_{pa} in 2007. As predicted last year, surveys indicate that cod recruitment is anticipated to be below the long-term mean both in 2009 and 2010, and also additionally in 2011.

Management objectives

A management plan has been implemented since 2004 (Annex 3.4.1) with the objectives of maintaining high long-term yield, year-to-year stability, and full utilization of all available information on stock dynamics. The plan aims to maintain F at $F_{pa} = 0.40$ and restrict between-year TAC change to $\pm 10\%$ unless SSB falls below B_{pa} , in which case the target F should be reduced.

Based on evaluations made in 2006 and 2007, ICES considers the management plan to be in accordance with the precautionary approach. If conditions change to outside the range assumed in management plan evaluation (with respect to biological conditions, assessment quality, and implementation error), the management plan may have to be revised.

Reference points

	Type	Value	Technical basis
Precautionary Approach	B_{lim}	220 000 t	change point regression
	B_{pa}	460 000 t	the lowest SSB estimate having >90% probability of remaining above B_{lim}
	F_{lim}	0.74	F corresponding to an equilibrium stock = B_{lim}
	F_{pa}	0.40	the highest F estimate having >90% probability of remaining below F_{lim}
Target	F_{pa}	0.40	

(unchanged since: 2003)

Yield and spawning biomass per Recruit F -reference points (2009):

	Fish Mort	Yield/R	SSB/R
	Ages 5-10		
Average last 3 years	0.41	1.12	1.93
F_{max}	0.25	1.16	3.76
$F_{0.1}$	0.13	1.07	7.11
F_{med}	0.80	0.99	0.65

HCR evaluation has shown that candidates for reference points which are consistent with high long-term yields and a low risk of depleting the productive potential of the stock can be found at F_{pa} .

Single-stock exploitation boundaries

ICES advises on the basis of the existing management plan which implies landings of 577 500 t in 2010.

Exploitation boundaries in relation to existing management plans

The agreed management plan implies landings of 577 500 t in 2010 (maximum 10% change in TAC from 2009). This projection includes all landings and therefore the TAC must account for any unreported landings.

Exploitation boundaries in relation to high long-term yield, low risk of depletion of production potential, and considering ecosystem effects

The current fishing mortality is in the range of F_s that are associated with high long-term yield and low risk of depleting the production potential.

Exploitation boundaries in relation to precautionary limits

The agreed management plan has been found to be consistent with the precautionary approach and is therefore the basis for the advice.

Short-term implications

Outlook for 2010:

For the forecast the F in 2009 is set equal to F in 2008.

Basis: $F(2009) = F_{2008} = 0.30$; $SSB(2010) = 1353$; landings (2009) = 522.

Rationale	Catch¹⁾ (2010)	Basis	F (2010)	SSB (2011)	%SSB change²⁾	% TAC change³⁾
Zero catch	0	$0 * F_{sq}$	0	2184	+62	-100
Agreed management Plan ⁴⁾	577.5	$0.95 * F_{sq}$	0.28	1655	+22	+10
<i>Status quo</i>	605	$1.00 * F_{sq}$	0.30	1631	+21	+15
Precautionary Limits	776	F_{pa}	0.40	1479	+9	+48

Weights in '000 t.

¹⁾ Catch are total catches without IUU landings. If this figure is taken as TAC, no implementation error is assumed.

²⁾ SSB 2011 relative to SSB 2010.

³⁾ Catch 2010 relative to TAC 2009.

⁴⁾ Forecast based on 10% TAC increase.

Management considerations

The TAC for 2009 was set above the catch corresponding to the agreed management plan. The earlier testing of the agreed management plan presumed that the plan should be strictly followed for setting TAC, and the deviation from the management plan in last year is not considered to be a precautionary practice. ICES considers that application of the agreed management plan in 2010 has long-term benefits above the application of F_{pa} .

The estimates of unreported landings have been reduced considerably from 2006 to 2008, which can probably be attributed to the introduction of port state control in the NEAFC area from 1 May 2007. For 2008, the estimate of 15 000 t unreported landings is around 3% of the international reported catch.

Unreported landings will reduce the effect of management measures and will undermine the intended objectives of the harvest control rule. It is therefore important that management agencies ensure that all catches are counted against the TAC.

Management plan evaluations

This management plan was evaluated in 2007. No new evaluations were carried out this year.

Factors affecting the fisheries and the stock

Regulations and their effects

TAC regulations are in place but there is non-compliance, resulting in a significant amount of unreported landings in the past. The main mechanism used in avoiding quota control seems to be trans-shipping of fish from the Barents Sea.

There is growing evidence of discarding throughout the Barents Sea for most groundfish stocks. Discarding is illegal in Norway and Russia. Data on discarding are scarce, but attempts to obtain better quantification continue.

The fisheries are controlled by inspections of the trawler fleet at sea, i.e. by a requirement to report to catch control points when entering and leaving the EEZs, VMS satellite tracking for some fleets, and by random inspections of fishing vessels when landing the fish. Keeping a detailed fishing logbook on-board is mandatory for most vessels, and large parts of the fleet report to the authorities on a daily basis.

In addition to quotas, the fisheries are regulated by mesh size limitations, a minimum catching size, a maximum bycatch of undersized fish, maximum bycatch of non-target species, closure of areas with high densities of juveniles, and other seasonal and area restrictions. The effects of these regulations have not been evaluated.

Changes in fishing technology and fishing patterns

Since January 1997, sorting grids have been mandatory for the trawl fisheries in most of the Barents Sea and Svalbard area.

Impacts of the environment on the fish stock

The Northeast Arctic cod is characterized by significant year-to-year variations in the growth rate. In some years the mean weight of fish at the same age may differ by a factor of 2 or 3. Among the factors influencing cod growth are water temperature, food supply, and cod population abundance. Since 2008, the recruitment predictions include information on environmental drivers.

Northeast Arctic cod is an important predator on other species in the ecosystem, notably capelin. The management of Northeast Arctic cod will therefore have implications on the dynamics of these stocks. Changes in growth, maturity, and cannibalism are linked to the abundance of capelin. This linkage appears to be less pronounced in the recent period compared to the 1980s and 1990s. In recent years, maturation, growth, and cannibalism have been fairly stable in spite of the variation in the capelin stock.

Annual consumption of cod by seals and whales may be inversely related to capelin abundance.

Scientific basis

Data and methods

Analytical assessment is based on catch-at-age data, using one commercial cpue series and three survey series. The total effect of discarding is still unclear and requires more work before it can be included in the assessments. Estimates of cannibalism are included in the natural mortality. Two series of IUU catch were made available to ICES for the years 2002-2008, but the advice is based on one series only. Since 2008, the recruitment predictions have included information on environmental drivers (ice coverage, temperature and oxygen saturation at the Kola section, air temperature at Murman coast, and capelin biomass).

Information from fishing industry

Several Norwegian fishing vessels (13 oceanic and 21 coastal) provide regular sampling data for length and age. These data are used for estimating catch at age for the corresponding fleets. Russian fishing vessels with observers onboard provide similar information on catch length distribution and sample fish to receive data on length-age matrices.

Uncertainties in assessment and forecast

The main uncertainties in this assessment derive from the biased catch statistics and the inconsistencies in the surveys. Bias in the catch statistics appears to have decreased in recent years. The survey results from the two last years are not consistent with the results from the previous years. Some of this inconsistency may be explained by inadequate spatial coverage of surveys in 2006/2007.

Environmental conditions

Environmental drivers were used in estimating recruitment and temperature used for estimating cod consumption.

Comparison with previous assessment and advice

Compared to last year's assessment, the current assessment estimates the SSB in 2008 to be 18% higher and the mean F in 2007 to be 13 % lower. The method to forecast recruitment that changed last year has been used again this year. Advice is, as last year, based on the management plan.

Source of information

ICES. 2009. Report of the Arctic Fisheries Working Group, 21–27 April 2009. ICES CM 2009/ACOM:02.

Year	ICES Advice	Single-stock exploitation boundaries	Predicted catch corresp. to advice	Predicted catch corresp. to single-stock exploitation boundaries	Agreed TAC	Official landings	ICES landings	Unreported landings (included in ICES landings)
1987	Gradual reduction in F		595		560	552	523	
1988	F = 0.51; TAC (Advice November 87, revised advice May 88)		530 (320–360)		590 451	459	435	
1989	Large reduction in F		335		300	348	332	
1990	F at F_{low} ; TAC		172		160	210	212	25
1991	F at F_{low} ; TAC		215		215	294	319	50
1992	Within safe biological limits		250		356	421	513	130
1993	Healthy stock		256		500	575	582	50
1994	No long-term gains in increased F		649		700	795	771	25
1995	No long-term gains in increased F		681		700	763	740	
1996	No long-term gains in increased F		746		700	759	732	
1997	Well below F_{med}		< 993		850	792	762	
1998	F less than F_{med}		514		654	615	593	
1999	Reduce F to below F_{pa}		360		480	506	485	
2000	Increase B above B_{pa} in 2001		110		390		415	
2001	High prob. of $SSB > B_{pa}$ in 2003		263		395		426	
2002	Reduce F to well below 0.25		181		395		535	90
2003	Reduce F to below F_{pa}		305		395		552	115
2004	Reduce F to below F_{pa}		398		486		606	117
2005	Take into account coastal cod and redfish by-catches	Apply catch rule	485		485		641	166
2006	Take into account coastal cod and redfish by-catches	Apply amended catch rule	471		471		538	67
2007	Take into account coastal cod and redfish by-catches	F_{pa}	309		424		487	41
2008	Take into account coastal cod and redfish by-catches	Apply catch rule	409		430		464	15
2009	Take into account coastal cod and redfish by-catches	Apply catch rule	473		525			
2010	Take into account coastal cod and redfish by-catches	Apply catch rule	577.5					

Weights in '000 tonnes.

Unreported landings 2002–2008 corrected according to Table 3.4.1.1.



Figure 3.4.1.1 Northeast Arctic cod (Subareas I and II). Landings, fishing mortality, recruitment, SSB






Figure 3.4.1.2 Northeast Arctic cod (Subareas I and II). Stock and recruitment, yield, and precautionary approach.




Figure 3.4.1.3 Northeast Arctic cod (Subareas I and II). Historical performance of the assessments

Table 3.4.1.1 Northeast Arctic COD. Total landings (t) by fishing areas and unreported landings.

Year	Subarea I	Division IIa	Division IIb	Unreported catches	Total catch
1961	409 694	153 019	220 508		783 221
1962	548 621	139 848	220 797		909 266
1963	547 469	117 100	111 768		776 337
1964	206 883	104 698	126 114		437 695
1965	241 489	100 011	103 430		444 983
1966	292 253	134 805	56 653		483 711
1967	322 798	128 747	121 060		572 605
1968	642 452	162 472	269 254		1 074 084
1969	679 373	255 599	262 254		1 197 226
1970	603 855	243 835	85 556		933 246
1971	312 505	319 623	56 920		689 048
1972	197 015	335 257	32 982		565 254
1973	492 716	211 762	88 207		792 685
1974	723 489	124 214	254 730		1 102 433
1975	561 701	120 276	147 400		829 377
1976	526 685	237 245	103 533		867 463
1977	538 231	257 073	109 997		905 301
1978	418 265	263 157	17 293		698 715
1979	195 166	235 449	9 923		440 538
1980	168 671	199 313	12 450		380 434
1981	137 033	245 167	16 837		399 037
1982	96 576	236 125	31 029		363 730
1983	64 803	200 279	24 910		289 992
1984	54 317	197 573	25 761		277 651
1985	112 605	173 559	21 756		307 920
1986	157 631	202 688	69 794		430 113
1987	146 106	245 387	131 578		523 071
1988	166 649	209 930	58 360		434 939
1989	164 512	149 360	18 609		332 481
1990	62 272	99 465	25 263	25 000	212 000
1991	70 970	156 966	41 222	50 000	319 158
1992	124 219	172 532	86 483	130 000	513 234
1993	195 771	269 383	66 457	50 000	581 611
1994	353 425	306 417	86 244	25 000	771 086
1995	251 448	317 585	170 966		739 999
1996	278 364	297 237	156 627		732 228
1997	273 376	326 689	162 338		762 403
1998	250 815	257 398	84 411		592 624
1999	159 021	216 898	108 991		484 910
2000	137 197	204 167	73 506		414 870
2001	142 628	185 890	97 953		426 471
2002 ²	184 789	189 013	71 242	90000/21716	535045/466760
2003 ²	163 109	222 052	51 829	115000/27748	551990/464738
2004 ²	177 888	219 261	92 296	117000/30000	606445/519445
2005 ²	159 573	194 644	121 059	166000/41000	641276/516276
2006 ²	159 851	204 603	104 743	127000/28000	596197/497197
2007 ²	152 522	195 383	97 891	41087/8757	486883/454553
2008 ¹	144905	203244	101022	15000/0	464171/449171

¹ Provisional figures.² two alternative estimates (see Chapter 3.1.3 of the 2008 AFWG Report for further details)

Table 3.4.1.2 Northeast Arctic COD. Total nominal landings ('000 t) by trawl and other gear for each area, data provided by Working Group members.

Year	Sub-area I		Division IIa		Division IIb	
	Trawl	Others	Trawl	Others	Trawl	Others
1967	238.0	84.8	38.7	90.0	121.1	-
1968	588.1	54.4	44.2	118.3	269.2	-
1969	633.5	45.9	119.7	135.9	262.3	-
1970	524.5	79.4	90.5	153.3	85.6	-
1971	253.1	59.4	74.5	245.1	56.9	-
1972	158.1	38.9	49.9	285.4	33.0	-
1973	459.0	33.7	39.4	172.4	88.2	-
1974	677.0	46.5	41.0	83.2	254.7	-
1975	526.3	35.4	33.7	86.6	147.4	-
1976	466.5	60.2	112.3	124.9	103.5	-
1977	471.5	66.7	100.9	156.2	110.0	-
1978	360.4	57.9	117.0	146.2	17.3	-
1979	161.5	33.7	114.9	120.5	8.1	-
1980	133.3	35.4	83.7	115.6	12.5	-
1981	91.5	45.1	77.2	167.9	17.2	-
1982	44.8	51.8	65.1	171.0	21.0	-
1983	36.6	28.2	56.6	143.7	24.9	-
1984	24.5	29.8	46.9	150.7	25.6	-
1985	72.4	40.2	60.7	112.8	21.5	-
1986	109.5	48.1	116.3	86.4	69.8	-
1987	126.3	19.8	167.9	77.5	129.9	1.7
1988	149.1	17.6	122.0	88.0	58.2	0.2
1989	144.4	19.5	68.9	81.2	19.1	0.1
1990	51.4	10.9	47.4	52.1	24.5	0.8
1991	58.9	12.1	73.0	84.0	40.0	1.2
1992	103.7	20.5	79.7	92.8	85.6	0.9
1993	165.1	30.7	155.5	113.9	66.3	0.2
1994	312.1	41.3	165.8	140.6	84.3	1.9
1995	218.1	33.3	174.3	143.3	160.3	10.7
1996	248.9	32.7	137.1	159.0	147.7	6.8
1997	235.6	37.7	150.5	176.2	154.7	7.6
1998	219.8	31.0	127.0	130.4	82.7	1.7
1999	133.3	25.7	101.9	115.0	107.2	1.8
2000	111.7	25.5	105.4	98.8	72.2	1.3
2001	119.1	23.5	83.1	102.8	95.4	2.5
2002	147.4	37.4	83.4	105.6	69.9	1.3
2003	146.0	17.1	107.8	114.2	50.1	1.8
2004	154.4	23.5	100.3	118.9	88.8	3.5
2005	132.4	27.2	87.0	107.7	115.4	5.6
2006	141.8	18.1	91.2	113.4	100.1	4.6
2007	129.6	22.9	84.8	110.6	91.6	6.3
2008 ¹	123.8	21.1	94.8	108.4	95.3	5.7

¹ Provisional figures.

Table 3.4.1.3 Northeast Arctic cod (Subareas I and II). Weights in t.

	Recruitment Age 3/thousands	SSB	Landings	Mean F ages 5-10
1946	728139	1112776	706000	0.1857
1947	425311	1165059	882017	0.3047
1948	442592	1019114	774295	0.3398
1949	468348	729879	800122	0.3619
1950	704908	615339	731982	0.3566
1951	1083753	568705	827180	0.3966
1952	1193111	520599	876795	0.5348
1953	1590377	396417	695546	0.3572
1954	641584	429694	826021	0.3879
1955	272778	346919	1147841	0.5437
1956	439602	299823	1343068	0.6401
1957	804781	207840	792557	0.5089
1958	496824	195377	769313	0.5169
1959	683690	432489	744607	0.5596
1960	789653	383479	622042	0.4789
1961	916842	404228	783221	0.6348
1962	728338	311678	909266	0.7576
1963	472064	208207	776337	0.9866
1964	338678	186570	437695	0.6789
1965	776941	102315	444930	0.5533
1966	1582560	120722	483711	0.5302
1967	1295416	129784	572605	0.5439
1968	164955	227215	1074084	0.5704
1969	112039	151870	1197226	0.8292
1970	197105	224482	933246	0.7493
1971	404774	311662	689048	0.5956
1972	1015319	346511	565254	0.6928
1973	1818949	332913	792685	0.6020
1974	523916	164491	1102433	0.5633
1975	621616	142028	829377	0.6595
1976	613942	171238	867463	0.6457
1977	348054	341385	905301	0.8379
1978	638490	241536	698715	0.9406
1979	198490	174699	440538	0.7264
1980	137735	108253	380434	0.7241
1981	150868	166926	399038	0.8632
1982	151830	326133	363730	0.7583
1983	166831	327181	289992	0.7560
1984	397831	251087	277651	0.9161
1985	523674	193856	307920	0.7038
1986	1038825	170729	430113	0.8649
1987	286344	121243	523071	0.9510
1988	204644	202589	434939	0.9743
1989	172782	234716	332481	0.6602
1990	242749	316418	212000	0.2710
1991	411784	704745	319158	0.3210
1992	721110	887561	513234	0.4550
1993	894891	775177	581611	0.5528
1994	810451	614868	771086	0.8678
1995	656938	528795	739999	0.7880
1996	437487	571581	732228	0.6987
1997	715761	588550	762403	1.0340
1998	845028	385854	592624	0.9173
1999	548930	292770	484910	0.9897
2000	608558	240387	414868	0.8540
2001	525622	355291	426471	0.7188
2002	454021	497911	535045	0.6751
2003	712993	549453	551990	0.5345
2004	305204	660319	606445	0.6778
2005	597565	619234	641276	0.7006
2006	510258	621033	537642	0.5805
2007	880265	692491	486883	0.3519
2008	796022	763193	464171	0.2989
2009	564000	1079210		
Average	609452	415072	651219	0.6301

Annex 3.4.1 Northeast Arctic Cod Management Agreement

At the 33rd meeting of the Joint Russian–Norwegian Fisheries Commission (JRNC) in November 2004, the following decision was made:

“The Parties agreed that the management strategies for cod and haddock should take into account the following:

*conditions for high long-term yield from the stocks
achievement of year-to-year stability in TACs
full utilization of all available information on stock development*

On this basis, the Parties determined the following decision rules for setting the annual fishing quota (TAC) for Northeast Arctic cod (NEA cod):

estimate the average TAC level for the coming 3 years based on F_{pa} . TAC for the next year will be set to this level as a starting value for the 3-year period.

the year after, the TAC calculation for the next 3 years is repeated based on the updated information about the stock development, however the TAC should not be changed by more than +/- 10% compared with the previous year's TAC.

if the spawning stock falls below B_{pa} , the procedure for establishing TAC should be based on a fishing mortality that is linearly reduced from F_{pa} at B_{pa} , to $F=0$ at SSB equal to zero. At SSB-levels below B_{pa} in any of the operational years (current year, a year before and 3 years of prediction) there should be no limitations on the year-to-year variations in TAC.¹”

¹ This quotation is taken from point 5.1 in the Protocol of the 33rd session of The Joint Norwegian–Russian Fishery Commission and translated from Norwegian to English. For an accurate interpretation, please consult the text in the official languages of the Commission (Norwegian and Russian).