

ECOREGION Iceland and East Greenland
STOCK Golden redfish (*Sebastes marinus*) in Subareas V, VI, XII, and XIV

Advice for 2013

ICES advises on the basis of precautionary considerations that catches should be no more than 40 000 t.

Stock status

F (Fishing Mortality)		
		2009–2011
MSY (F_{MSY})	?	Unknown
Precautionary approach (F_{pa}, F_{lim})	?	Unknown
SSB (Spawning-Stock Biomass)		
		2010–2012
MSY ($B_{trigger}$)	?	Unknown
Precautionary approach (B_{pa}, B_{lim})	✓	Full reproductive capacity
Qualitative evaluation	↗	Increasing in main area




Figure 2.4.7.1 Golden redfish (*Sebastes marinus*) in Subareas V, VI, XII, and XIV. Top left: Landings by area, Top right: Survey biomass (± 1 standard error) from for Icelandic waters (spring survey). Bottom: Survey biomass and abundance indices for East and West Greenland. Left: Total biomass index, including one standard error. Right: Total biomass index split into pre-fishery recruits (17–30 cm) and fishable redfish (>30 cm).

In Division Va the survey index (U) has been increasing since 2008 and is currently far above U_{pa} . In Division XIVb (East Greenland) survey indices of both pre-fishery recruits and fishable size have increased in recent years. In Division Vb the Faroese groundfish survey indicates that the abundance has been decreasing since 2001.

Management plans

The regulation is based on TAC in Iceland and in Greenland, but through an effort system in the Faroe Islands. The separation of golden redfish and Icelandic slope *S. mentella* in the quota was implemented in the 2010/2011 fishing season. The TAC in Greenland is set for redfish, with no distinction being made between *S. marinus* and *S. mentella*.

Biology

Sebastes marinus is a species with late maturation (matures between 10 and 14 years old) and slow growth (can get older than 50 years) and is hence considered to be vulnerable to overexploitation. It can therefore only sustain low exploitation and management should be based on that consideration.

The fisheries

The majority of the golden redfish catch is taken in ICES Division Va, which has contributed 95–98% of the total landings since 1990. Between 90 and 95% of the golden redfish catch in Division Va is taken by bottom trawlers targeting redfish. The remaining catches are caught as bycatch in gillnet, longline, and *Nephrops* fishery. Average annual landings 2000–2011 have been 40 000 tonnes. *S. marinus* in Division Va is to a small extent caught in a mixed fishery with *S. mentella* (Icelandic slope).

Catch distribution	Total landings (2011) = 44.8 kt, where 94% was taken by bottom trawls and 6% by other gear-types. Discards considered very small.
---------------------------	---

Quality considerations

Due to the aggregating behavior of the species, survey indices are often largely composed of a few large hauls. This causes high CVs in the indices and large interannual fluctuation in estimates of biomass.

Scientific basis

Assessment type	Trends-based assessment.
Input data	Two survey indices (Icelandic spring survey since 1985 and autumn survey since 1996).
Discards and bycatch	Not incorporated in the assessment, but very small.
Indicators	Gadget model used as trend indicator.
Other information	Faroese groundfish surveys since 1994 and 1996, German groundfish survey on the Greenland shelf since 1985. Benchmarked in February 2012, further review of the model settings needed before 2013.
Working group report	NWWG

ECOREGION Iceland, East Greenland, Faroe Islands
STOCK Golden redfish (*Sebastes marinus*) in Subareas V, VI, XII, and XIV

Reference points

ICES suggests that the relative state of the stock be assessed through a survey biomass index series (U) in Icelandic waters.

	Type	Value	Technical basis
MSY Approach	MSY B_{trigger}	Undefined	
	F_{MSY}	Undefined	
Precautionary approach	U_{lim}	55	20% of highest observed survey index*.
	U_{pa}	155	60% of highest observed survey index*.
	F_{lim}	Undefined	
	F_{pa}	Undefined	

(unchanged since 1998)

* Technical basis for the survey index:

The basis for the calculation of the U_{pa} is the Icelandic spring groundfish survey index series starting in 1985. Since 1990 the average U has been around half of U_{max} – the highest observed index in the time-series (276 in 1987). This has not resulted in any strong year classes compared to higher U's. A precautionary U_{pa} is therefore proposed at $U_{\text{max}} * 0.6$, corresponding to the U's associated with the most recent strong year class. U is regarded as a proxy for SSB but represents the fishable biomass.

Outlook for 2013

No analytical assessment can be presented for this stock, therefore, fishing possibilities cannot be projected.

Precautionary approach

The new data (landings and surveys) suggest the stock is increasing. The stock seems to have increased, with catches around 40 000 t since 1995. ICES advises that catches in 2013 should be no more than 40 000 t.

Additional considerations

Management considerations

Sebastes marinus is a species with late maturation and slow growth and is hence considered to be vulnerable to overexploitation. It can therefore only sustain low exploitation and management should be based on that consideration.

The strong 1990 year class has been in the Icelandic fishery for a decade and will also sustain the stock in the short term. The 1996–2001 year classes are above average and have been recruiting to the fishery since 2006.

An exploratory assessment based on the Gadget model indicates that the fishing mortality has decreased since the early 1990s and is currently close to F_{max} (Figure 2.4.7.2).

Subarea XIV is an important nursery area for *S. mentella* and *S. marinus*. The survey index of the fishable stock of *S. marinus* in Subarea XIV has increased in recent years, but with a large measurement uncertainty (Figure 2.4.7.2). Measures to protect juvenile redfish in Subarea XIV should be continued (sorting grids in the shrimp fishery).

In Subarea XIV redfish and cod are found in the same areas and depths and historically these species have been taken in the same fisheries. For 2012, ICES advises that no fishery should take place on cod in Greenland waters. Management measures should be put in place that minimize catches of cod in a directed fishery for *S. marinus*.

No formal agreement on the management of *S. marinus* exists among the three coastal states, Greenland, Iceland, and the Faroe Islands. In Greenland and Iceland, the fishery is regulated by a TAC and in the Faroe Islands by effort limitation.

On average, about 5% of the total landings have been taken in Division Vb and Subareas VI and XIV. In 2009 a fishery targeting redfish was initiated in Subarea XIV. Total catches were 1118 t in 2009, 8266 t in 2010, and 8381 t in 2011. The fishery does not distinguish between species, but based on survey information, *S. marinus* is estimated to account for 20% of catches, i.e. 224 t in 2009, 1653 t in 2010, and 1676 t in 2011.

Regulations and their effects

A quick closure system was implemented in 1977 in Iceland to protect juvenile redfish. If more than 20% of a catch observed on board is below 33 cm a small area can be closed for at least two weeks. For this reason there is no minimum landing size for golden redfish. The effect of the quick closure has not been evaluated and since 2001 there have been relatively few quick closures on small golden redfish, or on average three every year. The reason for the few quick closures on small golden redfish is because large areas southwest and west of Iceland are closed permanently or temporarily for trawling to protect juvenile golden redfish. These areas were closed partly because of frequent quick closures on redfish fisheries in 1991–1995. The effects of these closed areas have not been evaluated, but the increase in the spring survey index since 2003 is partly related to increased aggregation of golden redfish in these areas.

Since the late 1980s in Division Va and since 2002 in Subarea XIV it has been mandatory in the shrimp fishery to use sorting grids in order to reduce bycatches of juvenile redfish in the shrimp fishery.

Uncertainty in the assessment

A single abundance index that covers the whole distributional range of the stock is not available. The exploratory assessment is based on a survey index from Division Va only and landings from all three areas. This approach may create a bias in the assessment that cannot be quantified.

Comparison with previous assessment and advice

The basis and the advice has not changed compared to last year.

Sources

ICES 2012a. Report of the Benchmark Workshop on Redfish Stocks, 1–8 February 2012. ICES CM 2012/ACOM:48.
ICES. 2012b. Report of the North-Western Working Group, 26 April–3 May 2012. ICES CM 2012/ACOM:07.




Figure 2.4.7.2 Golden redfish in Subareas V, VI, XII, and XIV. Summary of stock assessment (weights in thousand tonnes.).




Figure 2.4.7.3 Golden redfish (*Sebastes marinus*) in Subareas V, VI, XII, and XIV. CpuE in the Faroese spring groundfish survey and the summer groundfish survey in ICES Division Vb.

Table 2.4.7.1 Golden redfish (*Sebastes marinus*) in Subareas V, VI, XII, and XIV. ICES advice, management, and landings.

Year	ICES Advice	Predicted catch corresp. to advice	Iceland TAC ^{1,6}	Greenland TAC ⁷	<i>S. marinus</i> ICES landings
1987	No increase in F	83	95		77
1988	No increase in F	84	85		90
1989	TAC ¹	117 ¹	77		57
1990	TAC ¹	116 ¹	80		67
1991	Precautionary TAC	77 (117 ¹)	55 ⁵		56
1992	Precautionary TAC	76 (116 ¹)	90		56
1993	Precautionary TAC ¹	120 ¹	104		50
1994	Precautionary TAC, if required	100 ¹	90		43
1995	TAC	90 ¹	77		45
1996	TAC for Division Va (28); precautionary TAC for Division Vb and Subarea XIV (4)	32 ²	65		37
1997	Effort 75% of 1995 value	32 ²	65		40
1998	Effort reduced in steps of 25% from the 1995 level	37.2 ²	65		39
1999	Effort not increased compared to 1997	35 ²	65		42
2000	Catch not increased compared to 1998	35 ²	60		44
2001	Effort not increased compared to 1999	33 ^{2,3}	57		37
2002	25% reduction in effort	29 ⁴	65		51
2003	25% reduction in effort(2001)	31 ⁴	60		39
2004	25% reduction in effort(2002)	37.4 ⁴	57		33.4
2005	Maintain fishable biomass above U _{pa}	37 ⁴	57		45.4
2006	Maintain fishable biomass above U _{pa}	37 ⁴	57		42.2
2007	Maintain fishable biomass above U _{pa}	37 ⁴	57	5	39.1
2008	Maintain fishable biomass above U _{pa}	37 ⁴	57	1	46.3
2009	Maintain fishable biomass above U _{pa}	< 30	50		39.2
2010	Maintain fishable biomass above U _{pa}	< 30	50	6	38.7
2011	Same advice as last year	< 30	37.5	8	44.8
2012	Maintain catches	< 40	40	8	
2013	Maintain catches	< 40			

Weights in '000 t.

¹ Deep-sea *S. mentella* and *S. marinus* combined.

² *S. marinus* only.

³ In Division Va only.

⁴ Both Divisions Va and Vb and Subarea XIV.

⁵ Year ending 31 August.

⁶ From 1992 onwards: Quota year September–August

⁷ Demersal redfish (*Sebastes marinus* and *S. mentella*).

Table 2.4.7.2 Golden redfish (*Sebastes marinus*) in Subareas V, VI, XII, and XIV. Official landings (in tonnes) by area.

Year	Area				Total
	Va	Vb	VI	XIV	
1978	31,300	2,039	313	15,477	49,129
1979	56,616	4,805	6	15,787	77,214
1980	62,052	4,920	2	22,203	89,177
1981	75,828	2,538	3	23,608	101,977
1982	97,899	1,810	28	30,692	130,429
1983	87,412	3,394	60	15,636	106,502
1984	84,766	6,228	86	5,040	96,120
1985	67,312	9,194	245	2,117	78,868
1986	67,772	6,300	288	2,988	77,348
1987	69,212	6,143	576	1,196	77,127
1988	80,472	5,020	533	3,964	89,989
1989	51,852	4,140	373	685	57,050
1990	63,156	2,407	382	687	66,632
1991	49,677	2,140	292	4,255	56,364
1992	51,464	3,460	40	746	55,710
1993	45,890	2,621	101	1,738	50,350
1994	38,669	2,274	129	1,443	42,515
1995	41,516	2,581	606	62	44,765
1996	33,558	2,316	664	59	36,597
1997	36,342	2,839	542	37	39,761
1998	36,771	2,565	379	109	39,825
1999	39,824	1,436	773	7	42,040
2000	41,187	1,498	776	89	43,550
2001	35,067	1,631	535	93	37,326
2002	48,570	1,941	392	189	51,092
2003	36,577	1,459	968	215	39,220
2004	31,686	1,139	519	107	33,451
2005	42,593	2,484	137	115	45,329
2006	41,521	656	0	34	42,211
2007	38,364	689	0	83	39,134
2008	45,538	569	64	80	46,251
2009	38,442	462	50	224	39,177
2010	36,155	620	220	1,653	38,648
2011 ¹⁾	42,605	493	83	1,676	44,875

¹⁾ Provisional.