

Short-term changes in abundance and population structure of dominant pelagic chaetognaths in the Oyashio region during the spring phytoplankton bloom

Yoshiyuki Abe¹, Hiroomi Miyamoto², Kohei Matsuno³, Atsushi Yamaguchi¹ and Ichiro Imai¹

In the Oyashio region, dominant water masses are switched at the surface layer within a short period during spring. Simultaneously, a large phytoplankton bloom is known to occur at the surface layer, and nearly half of the annual primary production is concentrated during spring. These drastic changes in water mass and food condition are expected to strongly affect macrozooplankton population dynamics. However, their effects on chaetognath population remain unknown. To evaluate the effects of water mass exchange and the spring phytoplankton bloom, we analyzed short-term changes in the population structure, growth rate, gut contents and predation impact of the three dominant chaetognaths (*Eukrohnia hamata*, *Parasagitta elegans* and *Pseudosagitta scrippsae*) in the Oyashio region during March–April 2007. Eleven samples were collected by 0-200 m oblique tow of Bongo nets at night during March 9 to April 30. Effects of water mass exchange were significant for *E. hamata* and *P. elegans*. During the sampling period, significant growth was observed for two dominant species (*E. hamata* and *P. elegans*). Daily growth rate was 39-50 $\mu\text{m day}^{-1}$ for *E. hamata* and 42-101 $\mu\text{m day}^{-1}$ for *P. elegans*. Mean predation impact of *P. elegans* at 0-200 m was 0.194 no. prey consumed $\text{m}^{-3} \text{day}^{-1}$, and that of *P. scrippsae* was 0.028 no. prey consumed $\text{m}^{-3} \text{day}^{-1}$. These values corresponded with 0-0.097% (*P. elegans*) or 0-0.043% (*P. scrippsae*) of the total zooplankton abundance during the phytoplankton bloom.

Keywords: spring bloom, chaetognaths, water mass, growth, predation impact

¹ Graduate School of Fisheries Science, Hokkaido University, Hakodate, Japan. E-mail: y.abe@fish.hokudai.ac.jp

² Tohoku National Fisheries Research institute, Fisheries Research Agency, Shiogama, Japan.

³ National Institute of Polar Research, Tachikawa, Japan.