

7.1* Bay of Biscay and the Iberian Coast Ecoregion – Ecosystem overview

Ecoregion description

The ICES Bay of Biscay and the Iberian Coast ecoregion covers the southwestern shelf seas and adjacent deeper eastern Atlantic Ocean waters of the EU. The ecoregion includes waters from Brittany to the Gulf of Cadiz; four key areas constitute the ecoregion:

- the Bay of Biscay, characterized by a wide shelf extending west of France. Upwelling events occur in summer off southern Brittany and low-salinity water lenses are associated with the river outflows of the Landes coastline;
- the northern Iberian Shelf, characterized by a narrow shelf with summer upwelling events off Galicia;
- the western Iberian Shelf, characterized by a narrow shelf west of Portugal with summer upwelling events;
- the Gulf of Cadiz, characterized by a wider shelf strongly influenced by input of warm Mediterranean waters.

The ecoregion includes parts of three Exclusive Economic Zones (EEZs) of EU Member States and a small portion of high seas. Fisheries in the Bay of Biscay and the Iberian Coast are managed under the Common Fisheries Policy (CFP), with some fisheries managed by the North East Atlantic Fisheries Commission (NEAFC) and by coastal states. Responsibility for salmon fishery management lies with the North Atlantic Salmon Conservation Organization (NASCO) and for large pelagic fish with the International Commission for the Conservation of Atlantic Tunas (ICCAT). Fisheries advice is provided by the International Council for the Exploration of the Sea (ICES), the European Commission's Scientific Technical and Economic Committee for Fisheries (STECF), and the South West Waters Advisory Council (SWWAC). Environmental policy is managed by national agencies and OSPAR, with advice being provided by national agencies, OSPAR, the European Environment Agency (EEA), and ICES. International shipping is managed under the International Maritime Organization (IMO).




Figure 7.1.1[†] The Bay of Biscay and the Iberian Coast ecoregion, showing EEZs and larger offshore Natura 2000 sites.

* Version2 ; section number corrected

[†] Version 2; figure number corrected




Figure 7.1.2[‡] Catchment area for the Bay of Biscay and the Iberian Coast ecoregion, showing major cities, ports, and ICES areas.

Key signals within the environment and the ecosystem

This ecoregion is characterized by marked seasonal wind-driven upwellings (see above) that strongly affect the productivity of the system.

- The seasonal pattern can be disrupted by (i) the Mediterranean water, which spills over into the Atlantic through the Strait of Gibraltar in the Gulf of Cadiz subregion, and by (ii) freshwater discharges from main rivers.
- Large-scale meteorological pressure differences over the North Atlantic can cause speed and direction anomalies of the wind during winter that have been linked with other parameters, with the recruitment of commercially important species such as anchovy *Engraulis encrasicolus*, sardine *Sardina pilchardus*, and horse mackerel *Trachurus trachurus*.
- The production of species such as sardine, southern hake *Merluccius merluccius*, and Norway lobster *Nephrops norvegicus* varies with the North Atlantic Oscillation meteorological pressure differences, with a time lag of *circa* three years.
- Fishing effort has been reduced in the Bay of Biscay and Iberian coast since the 2002 CFP reforms. Some stocks (i.e. anglerfish *Lophius* spp. and anchovy) are now fished at or below maximum sustainable yield (MSY) fishing mortality targets (F_{MSY}) or with spawning-stock biomass (SSB) above reference points; some others are still above target (i.e. hake *Merluccius merluccius* and megrim *Lepidorhombus* spp.), but for most stocks there is insufficient information to define reference points.

Pressures

The five most important pressures in the Bay of Biscay and Iberian Coast ecoregion are the selective extraction of species, abrasion, smothering, substrate loss, and nutrient and organic enrichment. These pressures are linked mainly to the following human activities: fishing, aquaculture, coastal construction, land-based industry, maritime transport, agriculture, dredging, and

[‡] Version 2; figure number corrected

offshore structures (Figure 7.1.3). Other pressures include the introduction of contaminating compounds, introduction of non-indigenous species, and underwater sound.

The pressures are described in the ICES glossary of pressures

http://www.ices.dk/sites/pub/Publication%20Reports/Advice/2016/2016/Pressure_Glossary_for_ecosystem_overviews.pdf.




Figure 7.1.3[§] Bay of Biscay and Iberian Coast ecoregion overview with the major regional pressures, human activities, and state of the ecosystem components. The width of lines indicates the relative importance of individual links.

Selective extraction of species

Fishing is the main activity contributing to this pressure in the Bay of Biscay and the Iberian Coast ecoregion. Both demersal and pelagic commercial fisheries occur in most parts of the ecoregion. Recreational fishery is becoming a relatively important activity and is in some cases taken into consideration for the management of marine fisheries. Tourism is also linked to aquatic and marine activities that contribute to the increase of this pressure in coastal areas. This pressure has four main effects on the ecosystem and its components, described below.

Impacts on commercial stocks

Figure 7.1.4 shows the historical evolution of fishing mortality and spawning-stock biomass relative to reference points by fish guild in the Bay of Biscay and the Iberian Coast ecoregion. A general decrease of fishing effort in the region (in many cases through reduction of the fleet) has contributed to an overall decline in the fishing mortality (F) of commercial fish stocks since

[§] Version 2; figure number corrected

1988. The mean F is now closer to the level that produces maximum sustainable yield (MSY); as a consequence an increase in the mean spawning-stock biomass has been observed since 2002.

Stocks of small pelagics like sardine and anchovy are highly influenced by natural recruitment variability and are therefore prone to periodic collapses linked to oceanographic variability. These stocks are closely monitored and regulated by strict management.

The conservation (sustainability) status of cephalopod populations varies depending on the particular subregion with declining catch per unit effort (CPUE) of octopus species in Galicia and long-finned squid *Loligo forbesi* off western Portugal, and increasing CPUE of octopus species in western Portugal and squid species in the southern Bay of Biscay.




Figure 7.1.4** Time-series of average of relative fishing mortality (F to F_{MSY} ratio) and biomass (SSB to B_{MSY} trigger ratio) by fish guild. Mean F and mean SSB is by total number of stocks with reference points.

Impact on threatened and declining fish species

Stocks of several fish species have been adversely affected by fishing and are now on the OSPAR list of threatened and declining species (see full list below). These include the sturgeon *Acipenser sturi*, European eel *Anguilla anguilla*, gulper shark *Centrophorus granulosus*, skates and rays like *Dipturus batis*, *Raja montagui*, and *Rostroraja alba*, spurdog *Squalus acanthias*, and salmon *Salmo salar*. Although there are no TACs for these species and some are prohibited to be landed under EU law, several species are vulnerable to existing fisheries. Common skates, and less often spurdogs, are caught as bycatch in demersal trawl fisheries while deepwater sharks are caught in the mixed deep-water trawl fishery.

Impacts on foodwebs

Fishing can disturb the foodweb. Predator–prey relationships can change, depending on the species and on the amount of food (prey) that is available for a given predator. Poor management of fishing for one species could have an adverse effect on the

** Version 2; figure number corrected

whole foodweb. Multispecies assessment methods can account for some of these interactions and guide appropriate management measures.

Indicators like the large fish indicator (LFI) index (describing the proportion – by weight – of the demersal fish community on survey catch larger than regional length thresholds) can be used to monitor changes in the fish populations. In the Bay of Biscay, the LFI index has shown a positive temporal trend since the year 2000 (Figure 7.1.5).




Figure 7.1.5^{††} Time-series of the large fish indicator (LFI) in the Bay of Biscay (ICES, 2013a).

There is no trend in the LFI in Portuguese waters. The index shows high interannual variability (Figure 7.1.6).




Figure 7.1.6^{††} Time-series of the large fish indicator (LFI) in Portuguese waters (ICES, 2013a).

Impacts on seabirds and marine mammals

Observation of marine mammal bycatch has occurred in certain fisheries off France and in a few off Galicia. Harbour porpoises *Phocoena phocoena* are being caught as bycatch off Iberia in set nets to the extent that the local population of the species may become extinct. Set net fisheries and pelagic trawls, particularly those for seabass *Dicentrarchus labrax*, have caught common dolphins *Delphis delphinus* and striped dolphins *Stenella coeruleoalba*. Seabird bycatch seems likely to be part of the reason for the loss of the Iberian form of the common guillemot *Uria aalge* and some other seabird species.

Abrasion

This pressure principally affects the seabed habitats and it is associated with bottom contacting mobile gear, in particular beam trawling, otter trawling, and local activities linked to tourism such as anchoring.

Using vessel monitoring system (VMS) and logbook data ICES estimates that mobile bottom trawls used by commercial fisheries in the 12 m+ vessel category have been deployed over approximately 99 000 km² of the ecoregion in 2013, corresponding to ca. 13.2% of the ecoregion’s spatial extent. Fishing is mainly concentrated along the shelf edge, largely in the Bay of Biscay and off the western Iberian coast (Figure 7.1.7 – note that the figure excludes Spanish fishing effort). Overall, there was a gradual decrease in the proportion of swept seafloor of ca. 2% between 2009 and 2013.

^{††} Version 2; figure number corrected

^{††} Version 2, figure number corrected




Figure 7.1.7⁵⁵ Surface and subsurface abrasion pressure expressed as the swept-area ratio obtained from VMS data from 2013 in the Bay of Biscay and Iberian Coast ecoregion. The figure excludes Spanish fishing effort. (ICES, 2015b.)

Smothering

Smothering is caused by several human activities in this ecoregion: extraction of aggregates (sand and gravel), disposal of materials on the seafloor, and navigational dredging for shipping as well as bottom trawling in soft sediment areas. The main dredging sites are found in France in the harbours and estuaries of the Loire (Nantes), Gironde (Bordeaux), and Adour (Bayonne), and in Spain (Avilés, Vilagarcía, and Huelva). This pressure affects both the benthic community structure and its productivity.

Substrate loss

Activities such as coastal construction, renewable energy devices, aquaculture, and fishing, all contribute to either sealing or changing natural substrates in this area.

Marine and coastal habitats have been lost in recent decades to land claim for port, industry, residential development and agriculture, coastal defences (including dykes, seawalls, and beach nourishment schemes), aquaculture infrastructure, shipping channels, roads, piers, marinas, and waste water treatment facilities.

Nutrient and organic enrichment

The input of nutrients is a relatively important pressure in coastal areas, particularly off areas of intensive agriculture and certain industries. Rivers account for most waterborne inputs of nitrogen and phosphorous. There is no clear trend in nitrate inputs in this region, in contrast to the decreasing trends in other EU waters. Maritime transport, international shipping, and aquaculture also contribute to this pressure.

Tourism and recreation along the coast is increasing in many parts of the ecoregion and seasonal management of the extra waste water produced is necessary. Eutrophication is mainly limited to coastal areas such as bays and estuaries with restricted circulation. Along the French coast, elevated levels of chlorophyll, nuisance phytoplankton species, and algal toxins have been observed.

⁵⁵ Version 2; figure number corrected

This pressure produces effects on the plankton community and on the overall productivity of the system.

Other pressures

Other relevant pressures in this ecoregion are:

- **Underwater noise**, caused by activities such as shipping, tourism and recreation, and renewable energy installations, which may affect marine mammals, fish, and other organisms using sound or pressure senses.
- **Introduction of non-indigenous species** happens primarily through shipping and aquaculture and may affect the benthic community and foodwebs.
- **Introduction of contaminating compounds**, due primarily to coastal discharges and maritime transport (shipping). This pressure can affect all ecosystem components but may accumulate in the foodweb, having an effect in particular on higher trophic levels (mammals and birds). Some of these compounds may be very stable and remain in the ecosystem for many decades after their introduction.

State of the ecosystem

Substrate and water masses

The substrate of the shelf of the Bay of Biscay and the Iberian Coast ecoregion is dominated sand and muddy-sand areas, with a large mud area in the Gulf of Cadiz (Figure 7.1.8).




Figure 7.1.8*** Major substrates on the shelf of the Bay of Biscay and Iberian coast (as compiled by EMODNET seabed habitats; www.emodnet-seabedhabitats.eu).

*** Version 2; figure number corrected

Pelagic habitats are dominated by the mixture of Mediterranean waters in the south and the influence of Atlantic waters in Portuguese, northern Spanish, and French waters. River run-off, particularly along the French coast, also influences the pelagic habitat.

Productivity

The most important features enhancing primary production and phytoplankton biomass are coastal upwelling, coastal run-off and river plumes, seasonal currents, and internal waves and tidal fronts. Concentrations of toxic dinoflagellates exhibit interannual variations determined mainly by changes in the upwelling regime, river run-off, inoculum size, and other environmental parameters.

Abundance of autotrophic picoplankton in the mid-shelf of the central part of the north Iberian coast do not show a significant long-term trend, but heterotrophic picoplankton show a significant long-term increase that has been associated to warming of the sea surface in this area. The zooplankton in the Bay of Biscay and Iberian Coast ecoregion do not show significant long-term changes.

Zooplankton

The most abundant zooplankton species are *Acartia* spp. and *Calanus helgolandicus*. Crab larvae are also important during winter months along the Portuguese coast. In recent years, copepod species such as *Temora stylifera*, characteristic of warmer waters, have appeared or increased in abundance (e.g.) related to increases in sea surface temperature. The seasonal cycle of zooplankton biomass is characterized by a bimodal pattern along the western Iberian coast, with peak biomass in April and August, caused by seasonal upwelling. Copepod abundance remains high throughout the year, with the highest abundances from August through November along the Iberian coast. The interannual variation of zooplankton abundance and biomass does not show clear trends in the ecoregion except for Galicia, where it couples with the upwelling cycles. There are differences in the structure of the zooplankton community along the north Iberian coast.

Cephalopods

Within this ecoregion the topographic diversity and the wide range of substrates result in many different habitats for cephalopods. In this region, the most abundant and commercially exploited species are Loliginidae (long-finned squid) and Sepiidae (cuttlefish). Abundance of Ommastrephidae (short-finned squid) increases westwards towards Galicia, and decreases to the south of the Iberian coast. Octopodidae are abundant and heavily exploited along the Iberian coast by a large artisanal fleet, with concomitant social relevance. There are indications of a decline in octopus biomass index in Galicia and an increase off western Portugal. Stocks of both long-finned squid and short-finned squid have declined in the southern Bay of Biscay.

Fish

Fish diversity is high in this ecoregion, reflecting its wide latitudinal dimension. The pelagic habitat is mainly dominated by sardine, anchovy, mackerel, horse mackerel, and blue-whiting *Micromesistius poutassou*. Some migratory species also appear in specific periods, such as tuna species (albacore *Thunnus alalunga* and bluefin *Thunnus thynnus*), which feed upon smaller pelagic fish. Hake is the most abundant predator species in the demersal community. Anglerfish, megrim, and sole are more abundant in the northern part of the ecoregion. Cold-water species such as whiting *Merlangius merlangus* and pollack *Pollachius pollachius* only occur north of Portugal. Skates, sharks, and deep-sea fish occur over the continental slope and in the deeper parts of this ecoregion. Trends in fishing pressure and stock size are presented in the 'Selective extraction of species' section.

Seabirds

The coasts of the Bay of Biscay and the western Iberian Peninsula are used by several seabird species for breeding. These include the European storm petrel *Hydrobates pelagicus*, European shag *Phalacrocorax aristotelis*, yellow-legged gull *Larus michahellis*, lesser black-backed gull *Larus fuscus*, black-legged kittiwake *Rissa tridactyla*, and common guillemot. Many more species use these waters for feeding in the non-breeding period. The most important species in terms of abundance are northern gannet *Morus bassanus*, gulls *Larus* spp. (seven species), Balearic shearwater *Puffinus mauretanicus*, Manx shearwater *Puffinus puffinus*, sooty shearwater *Puffinus griseus*, Cory's shearwater *Calonectris diomedea*, razorbill *Alca torda*,

and Atlantic puffin *Fratercula arctica*. Trends in the numbers of seabirds breeding around these seas are not known, with the exception of Iberian common guillemot and black-legged kittiwake that are either now extirpated or close to that state. Shags have also declined.

Marine mammals

No seals are common in the Bay of Biscay and Iberian Atlantic waters. Twelve species of cetacean occur commonly or are resident in the Celtic Seas: minke whale *Balaenoptera acutorostrata*, fin whale *Balaenoptera physalus*, harbour porpoise, short-beaked common dolphin *Delphinus delphis*, striped dolphin *Stenella coeruleoalba*, sperm whale *Physeter macrocephalus*, long-finned pilot whale *Globicephala melas*, northern bottlenose whale *Hyperoodon ampullatus*, Cuvier’s beaked whale *Ziphius cavirostris*, Sowerby’s beaked whale *Mesoplodon bidens*, Risso’s dolphin *Grampus griseus*, and bottlenose dolphin *Tursiops truncatus*. There is no information on trends in the abundance of marine mammals in this region.

Non-indigenous species

This region has 217 non-indigenous and cryptogenic (obscure or of unknown origin) species. The majority (55 species) arrived between 1950 and 1999. Since 2000, a total of 25 new species have been recorded of which 12 are new to Europe. Consequently, the annual rate of discovery of non-indigenous species increased from 1.1 per year during 1950–2014, to 1.7 per year during 2000–2014 (see Figure 7.1.9).

Shipping, in particular through ballast water and biofouling of hulls, as well as coastal water currents (secondary spread from neighbouring areas) are the main species introduction vectors, followed by aquaculture activities (predominantly by shellfish transfers).

Ecological impacts such as declines of native species and structural changes in the benthic communities have been observed. Other impacts include fouling of irrigation systems and clogging of fishing nets and aquaculture nets.




Figure 7.1.9⁺⁺⁺ Annual rate of new non-indigenous and cryptogenic species discoveries in the Bay of Biscay and Iberian Coast ecoregion during 1950–1999 and 2000–2014.

⁺⁺⁺ Version 2; figure number corrected

Threatened and declining species and habitats in the Bay of Biscay and Iberian Coast ecoregion

The threatened and declining species in the Bay of Biscay and Iberian Coast ecoregion according to OSPAR are shown in the table below.

SCIENTIFIC NAME	COMMON NAME
INVERTEBRATES	
<i>Nucella lapillus</i>	Dog whelk
SEABIRDS	
<i>Puffinus mauretanicus</i>	Balearic shearwater
<i>Sterna dougallii</i>	Roseate tern
<i>Uria aalge</i> – Iberian population (synonyms: <i>Uria aalge albionis</i> , <i>Uria aalge ibericus</i>)	Iberian guillemot
FISH	
<i>Acipenser sturio</i>	Sturgeon
<i>Alosa alosa</i>	Allis shad
<i>Anguilla anguilla</i>	European eel
<i>Centroscymnus coelolepis</i>	Portuguese dogfish
<i>Centrophorus granulosus</i>	Gulper shark
<i>Centrophorus squamosus</i>	Leafscale gulper shark
<i>Cetorhinus maximus</i>	Basking shark
<i>Dipturus batis</i> (synonym: <i>Raja batis</i>)	Common skate
<i>Raja montagui</i> (synonym: <i>Dipturus montagui</i>)	Spotted ray
<i>Hippocampus guttulatus</i> (synonym: <i>Hippocampus ramulosus</i>)	Long-snouted seahorse
<i>Hippocampus hippocampus</i>	Short-snouted seahorse
<i>Lamna nasus</i>	Porbeagle
<i>Petromyzon marinus</i>	Sea lamprey
<i>Rostroraja alba</i>	White skate
<i>Salmo salar</i>	Salmon
<i>Squalus acanthias</i>	[Northeast Atlantic] spurdog
<i>Squatina squatina</i>	Angel shark
REPTILES	
<i>Caretta caretta</i>	Loggerhead turtle
<i>Dermochelys coriacea</i>	Leatherback turtle
MARINE MAMMALS	
<i>Balaenoptera musculus</i>	Blue whale
<i>Eubalaena glacialis</i>	Northern right whale
<i>Phocoena phocoena</i>	Harbour porpoise

Threatened and declining habitats in the Bay of Biscay and Iberian Coast ecoregion according to OSPAR

HABITATS
Coral gardens
<i>Cymodocea</i> meadows
Deep-sea sponge aggregations
Intertidal mudflats
<i>Lophelia pertusa</i> reefs
<i>Modiolus modiolus</i> beds
<i>Ostrea edulis</i> beds
Seamounts
<i>Zostera</i> beds

Sources and acknowledgments

The content for the ICES regional ecosystem overviews is based on information and knowledge generated by the following ICES processes: Workshop on Benchmarking Integrated Ecosystem Assessment (WKBEMIA) 2012, ACOM/SCICOM Workshop on Ecosystem Overviews (WKECOVER) 2013, Workshop to draft advice on Ecosystem Overviews (WKDECOVER) 2013, and Advice drafting group to finalize draft Ecosystem Overviews (ADGECO) 2015, which provided the theoretical framework and final layout of the documents. The ICES integrated ecosystem assessment Working Group on Ecosystem Assessment of Western European Shelf Seas (WGEAWESS) contributed to the main sections of this overview. The following working groups contributed to draft the subsections on the state of the ecosystem components: Benthos Ecology Working Group (BEWG), Working Group on Multispecies Assessment Methods (WGSAM), Working Group on Zooplankton Ecology (WGZE), Working Group on Cephalopod Fisheries and Life History (WGCEPH), Working Group on Marine Mammal Ecology (WGMME), and Working Group on Introductions and Transfers of Marine Organisms (WGITMO).

The maps and GIS products have been produced by the ICES Secretariat using data from:

1. Exclusive Economic Zones. *Marineregions.org* (VLIZ)
2. Offshore Wind-farms. *OSPAR Commission*
3. Depth Contours. *General Bathymetric Chart of the Oceans (GEBCO)*
4. Natura 2000. *European Commission*
5. Ecoregions. *International Council for the Exploration of the Sea (ICES)*
6. Ports. *Global Shipping Lanes and Harbors (ESRI)*
7. Cities. *World Cities (ESRI)*
8. Rivers. *WISE Large Rivers and large lakes. European Environment Agency (EEA)*
9. ICES Areas. *International Council for the Exploration of the Sea (ICES)*
10. Catchment Area. *European Environment Agency (EEA). European Topic Centre on Inland, Coastal and Marine waters (ETC/ICM).*
11. Substrate maps. EU EMODNET seabed habitats; www.emodnet-seabedhabitats.eu
12. Non indigenous species. AquaNIS; <http://www.corpi.ku.lt/databases/index.php/aquanis>

Sources and references

Behrenfeld, M. J., and Falkowski, P. G. 1997. Photosynthetic rates derived from satellite-based chlorophyll concentration. *Limnology and Oceanography*, 42: 1–20.

Blanton, J. O., Atkinson, L. P., Fernandez de Castillejo, F., and Lavin, A. 1984. Coastal upwelling off the Rias Bajas, Galicia, northwest Spain I: hydrographic studies. *Rapports et Proces Verbaux des Reunions CIEM*, 183: 79–90.

- Borges, M. F., Santos, A. M. P., Crato, N., Mendes, H., and Mota, B. 2003. Sardine regime shifts off Portugal: A time series analysis of catches and wind conditions. *Scientia Marina*, 66(S1): 235–244.
- Borges, M. F., Mendes, H., and Santos, A. M. P. 2012. Sardine (*Sardina pilchardus*) recruitment is strongly affected by climate even at high spawning biomass in West Iberia/Canary upwelling system. *In Science and Management of Small Pelagics*. Edited by S. Garcia, M. Tandstad, and A. M. Caramelo. FAO Fisheries and Aquaculture Proceedings, No. 18: 237–244. Rome, FAO.
- Borja, A., Uriarte, A., Egana, J., Motos, L., and Valencia, V. 1998. Relationships between anchovy (*Engraulis encrasicolus*) recruitment and environment in the Bay of Biscay (1967–1996). *Fisheries Oceanography*, 7: 375–380.
- Botas, J. A., Fernández, E., Bode, A., and Anadón, R. 1990. A persistent upwelling off the Central Cantabrian Coast (Bay of Biscay). *Estuarine, Coastal and Shelf Science*, 30: 185–199.
- Cayan, D. R. 1992. Latent and sensible heat flux anomalies over the northern oceans: driving the sea surface temperature. *Journal of Physical Oceanography*, 22: 859–881.
- Fiúza, A. F. D., De Macedo, M. E., and Guerreiro, M. R. 1982. Climatological space and time-variation of the Portuguese coastal upwelling. *Oceanologica Acta*, 5: 31–40.
- Fraga, F. 1981. Upwelling off the Galician coast, northwest Spain. *In Coastal Upwelling*, pp. 176–182. Ed. by F. A. Richards, American Geophysical Union, Washington, D. C.
- Frid, C., Andonegi, E., Depestele, J., Judd, A., Rihan, D., Rogers, S. I., and Kenchington, E. 2012. The environmental interactions of tidal and wave energy generation devices. *Environmental Impact Assessment Review*, 32: 133–139.
- García-Lafuente, J., and Ruiz, J. 2007. The Gulf of Cádiz pelagic ecosystem. *Progress in Oceanography*, 74: 228–251.
- González-Pola, C., and Izquierdo, P. 2012. Estrategia Marina. Demarcación Marina Noratlántica. Parte IV: Descriptores del buen estado ambiental. Descriptor 7: Condiciones Hidrográficas. Evaluación inicial y buen estado ambiental. Ministerio de Agricultura, Alimentación y Medio Ambiente, Secretaría General Técnica, Centro de Publicaciones. NIPO: 208-12-175-8. <http://publicacionesoficiales.boe.es/>.
- González-Pola, C., Lavín, A., del Río, G. D., Cabanas, J., Ruiz-Villareal, M., Somavilla, R., Rodríguez, C., González-Nuevo, G., and Nogueira, E. 2012. Hidrografía y circulación. *In Cambio Climático y oceanografía en el Atlántico del Norte de España*, pages 69–98. Edited by A. Bode, A. Lavín, and L. Valdés. Temas de Oceanografía, Vol. 5, Instituto Español de Oceanografía, Ministerio de Ciencia e Innovación, ISBN 978-84-95877-08-6.
- ICES. 2011a. Report of the ICES Advisory Committee, 2011. ICES Advice 2011, Book 7. 118 pp.
- ICES. 2011b. Report of the Working Group on Anchovy and Sardine (WGANSA), 24–28 June 2011, Vigo, Spain. ICES CM 2011/ACOM:16. 8 pp.
- ICES. 2012a. Report of the Working Group on the Assessment of Southern Shelf Stocks of Hake, Monk and Megrim (WGHMM), 10–16 May 2012, ICES Headquarters, Copenhagen. ICES CM 2012/ACOM:11. 617 pp.
- ICES. 2012b. Report of the Working Group on Oceanic Hydrography (WGOH), 21–22 March 2012, ICES Headquarters, Copenhagen, Denmark. ICES CM 2012/SSGEF:03. 168 pp.
- ICES. 2012c. Report of the Workshop on Benchmarking Integrated Ecosystem Assessments (WKBEMIA), 27–29 November 2012, ICES Headquarters, Copenhagen, Denmark. ICES CM 2012/SSGRSP:08. 28 pp.
- ICES. 2013a. Report of the Workshop on DCF Indicators (WKIND), 21–25 October 2013, ICES Headquarters, Copenhagen, Denmark. ICES CM 2013/ACOM:38. 78 pp.
- ICES. 2013b. Report of the Working Group on the ICES ACOM/SCICOM Workshop on Ecosystem Overviews (WKECOVER), 7–11 January 2013, ICES HQ, Copenhagen, Denmark. ICES CM 2013/ACOM/SCICOM:01. 131 pp.
- ICES. 2013c. Report of the ICES Workshop to draft Advice on Ecosystem Overviews (WKDECOVER), 4–7 November, ICES HQ, Copenhagen. ICES CM 2013/ACOM/SCICOM:03. 11 pp.

ICES. 2013d. Report of the Working Group on Ecosystem Assessment of Western European Shelf Seas (WGEAWESS), 11–15 February 2013, Lisbon, Portugal. ICES CM 2013/SSGRSP:02. 164 pp.

ICES. 2015a. ICES glossary of principal pressures in ICES ecoregions.

http://www.ices.dk/sites/pub/Publication%20Reports/Advice/2016/2016/Pressure_Glossary_for_ecosystem_overviews.pdf.

ICES. 2015b. Indicators 5, 6, and 7 of DCF Annex XII. In Report of the ICES Advisory Committee, 2015. ICES Advice 2015, Book 1, Section 1.6.1.4.

ICES. 2015c. First and Second Interim Reports of the Working Group on Ecosystem Assessment of Western European Shelf Seas (WGEAWESS), 9–13 March 2015, 1. via WebEx and Correspondence 2. Cadiz, Spain. ICES CM 2015/SSGIEA:02. 32 pp.

ICES. 2015d. Interim Report of the Benthos Ecology Working Group (BEWG), 4–8 May 2015, Calvi, Corsica, France. ICES CM 2015/SSGEPD:10. 64 pp.

ICES. 2015e. Interim Report of the Working Group on Zooplankton Ecology (WGZE), 16–19 March 2015, Plymouth, UK. ICES CM 2015/SSGEPD:05. 44 pp.

ICES. 2015f. Report of the Working Group on Marine Mammal Ecology (WGMME), 9–12 February 2015, London, UK. ICES CM 2015/ACOM:25. 114 pp.

ICES. 2015g. Report of the Working Group on Introductions and Transfers of Marine Organisms (WGITMO), 18–20 March 2015, Bergen, Norway. ICES CM 2015/SSGEPI:10. 195 pp.

Koutsikopoulos, C., and Le Cann, B. 1996. Physical processes and hydrological structures related to the Bay of Biscay anchovy. *In* The European Anchovy and its environment. Edited by I. Palomera and P. Rubiés. Scientia Marina, 60 (S2): 9–19.

Lavín, A., Valdés, L., Gil, J., and Moral, M. 1998. Seasonal and interannual variability in properties of surface water off Santander, Bay of Biscay, 1991–1995. *Oceanologica Acta*, 21: 179–189.

Levitus, S., Antonov, J. I., Wang, J., Delworth, T. L., Dixon, K. W., and Broccoli, A. J. 2001. Anthropogenic warming of Earth's climate system. *Science*, 292(5515): 267–270.

Michel, S., Vandermeersch, F., and Lorange, P. 2009. Evolution of upper layer temperature in the Bay of Biscay during the last 40 years. *Aquatic Living Resources*, 22: 447–461.

Moita, T. 2001. Estrutura, variabilidade e dinâmica do Fitoplâncton na costa de Portugal Continental. PhD Thesis. University of Lisbon, Portugal. 272 pp.

Morán, X. A. G., López-Urrutia, A., and Calvo-Díaz, A. 2010. Increasing importance of small phytoplankton in a warmer ocean. *Global Change Biology*, 16: 1137–1144.

Morán, X. A. G., Bode, A., Calvo-Díaz, A., Díaz-Pérez, L., Suárez, L. A., Roura, A., Nogueira, E., and González-Nuevo, G. 2012. Capítulo 5: Picoplankton autotrófico y heterotrófico. *In* Cambio Climático y oceanografía en el Atlántico del Norte de España, pages 159–175. Edited by A. Bode, A. Lavín, and L. Valdés. *Temas de Oceanografía*, Vol. 5, Instituto Español de Oceanografía, Ministerio de Ciencia e Innovación, ISBN 978-84-95877-08-6.

Navarro, G., and Ruiz, J. 2006. Spatial and temporal variability of phytoplankton in the Gulf of Cádiz through remote sensing images. *Deep-Sea Research II*, 53: 1241–1260.

Nogueira, E., Pérez, F. F., and Ríos, A. F. 1997. Modelling thermohaline properties in an estuarine upwelling ecosystem (Ria de Vigo; NW Spain) using Box-Jenkins transfer function models. *Estuarine, Coastal and Shelf Science*, 44: 685–702.

ODEMM. Options for Delivering Ecosystem-Based Marine Management. <https://www.liverpool.ac.uk/odemmm/> and <http://www.odemm.com/>.

Ommer, R., Perry, I., Cochrane, K. L., and Cury, P. 2011. *World fisheries: a social-ecological analysis*. Wiley-Blackwell, London. 440 pp.

OSPAR. 2010. Quality Status Report 2010. OSPAR Commission, London. 176 pp.

Poulard, J. C., and Blanchard, F. 2005. The impact of climate change on the fish community structure of the eastern continental shelf of the Bay of Biscay. *ICES Journal of Marine Science*, 62: 1436–1443.

Preciado, I., Velasco, F., Valeiras, J., and Modica, L. 2012. Estrategia Marina. Demarcación Marina Noratlántica. Parte IV: Descriptores del buen estado ambiental. Descriptor 4: Redes Tróficas. Evaluación inicial y buen estado ambiental. Ministerio de Agricultura, Alimentación y Medio Ambiente, Secretaría General Técnica, Centro de Publicaciones. NIPO: 208-12-175-8. <http://publicacionesoficiales.boe.es/>.

Puillat, I., Lazure, P., Jegou, A. M., Lampert, L., and Miller, P. 2006. Mesoscale hydrological variability induced by northwesterly wind on the French continental shelf of the Bay of Biscay. *Scientia Marina*, 70(S1): 15–26.

Punzón, A., and Villamor, B. 2009. Does the timing of the spawning migration change the southern component of the Northeast Atlantic Mackerel (*Scomber scombrus*, L. 1758)? An approximation using fishery analyses. *Continental Shelf Research*, 29: 1195–1204.

Relvas, P., Luís, J., and Santos, A. M. P. 2009. Importance of the mesoscale in the decadal changes observed in the northern Canary upwelling system, *Geophysical Research Letters*, 36:L22601. doi:10.1029/2009GL040504.

Ruiz, J., Gonzalez-Quirós, R., Prieto, L., and Navarro, G. 2009. A Bayesian model for anchovy (*Engraulis encrasicolus*): the combined forcing of man and environment. *Fisheries Oceanography*, 18: 62–76.

Santos, A. M. P., Borges, M. F., and Groom, S. 2001. Sardine and horse mackerel recruitment and upwelling off Portugal. *ICES Journal of Marine Science*, 58: 589–596.

STECF. 2013. Scientific, Technical and Economic Committee for Fisheries (STECF) – Evaluation of Fishing Effort Regimes in European Waters – Part 2 (STECF-13-21). Publications Office of the European Union, Luxembourg, EUR 26327 EN, JRC86088. 863 pp.