

## Norway lobster (*Nephrops norvegicus*) in Division 6.a, Functional Unit 13 (West of Scotland, Firth of Clyde, and Sound of Jura)

### ICES advice on fishing opportunities


ICES advises that when the MSY approach is applied, and assuming that discard rates and fishery selection patterns do not change from the average of the years 2018–2020, catches in 2022 should be no more than 4235 tonnes (3607 tonnes for the Firth of Clyde and 628 tonnes for the Sound of Jura).

To ensure that the stock in Functional Unit (FU) 13 is exploited sustainably, management should be implemented at the FU level. In this particular FU, additional measures should be implemented to ensure that landings taken in each subarea (the Firth of Clyde and the Sound of Jura) are in line with the advice.

ICES notes the existence of a management plan, developed and adopted by one of the relevant management authorities for Subarea 6. ICES considers this plan to be precautionary when implemented at the FU level.

### Stock development over time

Fishing pressure on the stock is below  $F_{MSY}$  and stock size is above  $MSY B_{trigger}$ .



**Figure 1** Norway lobster in Division 6.a, Functional Unit 13. Summary of the stock assessment. Catches (discard data only available from 1990), harvest rate (sum of landings and dead discards in numbers, divided by stock abundance), and stock abundance (underwater TV survey). Harvest rates before 2006 may be underestimated because of the underreporting of landings. Harvest rates are calculated using the total catch divided by the stock abundance for the two subareas combined. For the Sound of Jura, an interpolated value for abundance in 2020 is used (average of 2019 and 2021).

## Catch scenarios

**Table 1** Norway lobster in Division 6.a, Functional Unit 13. The basis for the catch scenarios.

### Firth of Clyde

| Variable                          | Value | Notes  |
|-----------------------------------|-------|--|
| Stock abundance (2022)            | 1414  | UWTV Survey 2021; individuals in millions  |
| Mean weight in projected landings | 17.45 | Average 2018–2020 (combined for the Firth of Clyde and the Sound of Jura); in grammes                              |
| Mean weight in projected discards | 8.06  | Average 2018–2020 (combined for the Firth of Clyde and the Sound of Jura); in grammes                              |
| Projected discard rate            | 10.8  | Average 2018–2020 (combined for the Firth of Clyde and the Sound of Jura); percentage by number of the total catch |
| Discards survival rate            | 25    | Percentage by number of the discards   |

### Sound of Jura

| Variable                          | Value | Notes  |
|-----------------------------------|-------|--|
| Stock abundance (2022)            | 310   | UWTV Survey 2021; individuals in millions  |
| Mean weight in projected landings | 17.45 | Average 2018–2020 (combined for the Firth of Clyde and the Sound of Jura); in grammes                              |
| Mean weight in projected discards | 8.06  | Average 2018–2020 (combined for the Firth of Clyde and the Sound of Jura); in grammes                              |
| Projected discard rate            | 10.8  | Average 2018–2020 (combined for the Firth of Clyde and the Sound of Jura); percentage by number of the total catch |
| Discards survival rate            | 25    | Percentage by number of the discards   |

**Table 2** Norway lobster in Division 6.a, Functional Unit 13. Annual catch scenarios. All weights are in tonnes. The figures in the table are rounded. Calculations were done with unrounded inputs, and computed values may not match exactly when calculated using the rounded figures in the table. Catch scenarios assuming discarding continues at the recent average rate.

### Firth of Clyde

| Basis                      | Total catch    | Dead removals | Projected landings | Projected dead discards | Projected surviving discards | % harvest rate* | % advice change** |
|----------------------------|----------------|---------------|--------------------|-------------------------|------------------------------|-----------------|-------------------|
|                            | PL + PDD + PSD | PL + PDD      | PL                 | PDD                     | PSD                          | for PL + PDD    |                   |
| ICES advice basis          |                |               |                    |                         |                              |                 |                   |
| MSY approach               | 3607           | 3559          | 3416               | 143                     | 48                           | 15.1            | -25               |
| Other scenarios            |                |               |                    |                         |                              |                 |                   |
| F <sub>MSY lower</sub>     | 2364           | 2333          | 2239               | 94                      | 31                           | 9.9             | -51               |
| F <sub>MSY upper</sub> *** | 3607           | 3559          | 3416               | 143                     | 48                           | 15.1            | -25               |
| F <sub>2020</sub>          | 2245           | 2215          | 2126               | 89                      | 30                           | 9.4             | -53               |

### Sound of Jura

| Basis                      | Total catch    | Dead removals | Projected landings | Projected dead discards | Projected surviving discards | % harvest rate* | % advice change** |
|----------------------------|----------------|---------------|--------------------|-------------------------|------------------------------|-----------------|-------------------|
|                            | PL + PDD + PSD | PL + PDD      | PL                 | PDD                     | PSD                          | for PL + PDD    |                   |
| ICES advice basis          |                |               |                    |                         |                              |                 |                   |
| MSY approach               | 628            | 620           | 595                | 25                      | 8                            | 12.0            | -0.95             |
| Other scenarios            |                |               |                    |                         |                              |                 |                   |
| F <sub>MSY lower</sub>     | 493            | 486           | 466                | 20                      | 7                            | 9.4             | -22               |
| F <sub>MSY upper</sub> *** | 628            | 620           | 595                | 25                      | 8                            | 12.0            | -0.95             |
| F <sub>2020</sub>          | 493            | 486           | 466                | 20                      | 7                            | 9.4             | -22               |

\* By number.

\*\* Advice values for 2022 are relative to the 2021 advice (MAP F<sub>MSY</sub> advice of 4791 tonnes for the Firth of Clyde, and 634 tonnes for the Sound of Jura).

\*\*\* F<sub>MSY upper</sub> = F<sub>MSY</sub> for this stock.

The advice for 2022 is lower than for 2021 because of lower observed stock abundances.

### Basis of the advice

**Table 3** Norway lobster in Division 6.a, Functional Unit 13. The basis of the advice.

| Advice basis    | MSY approach  |
|-----------------|---|
| Management plan | ICES is aware of the EU multiannual management plan (MAP) that has been agreed for this stock (EU, 2019) and considers it to be precautionary when implemented at the FU level. There is no agreement with UK regarding this plan, and it is not used as the basis of the advice for this stock. ICES provides catch scenarios consistent with the $F_{MSY}$ ranges in the MAP. |

### Quality of the assessment

Discard sampling in 2020 was impacted by the COVID-19 pandemic, with no samples collected in FU 13. Estimates of discard rates for all quarters in the assessment were based on mean discard rates across all quarters 2017–2019 (ICES, 2021a). Landings in quarter 2 were not sampled due to COVID-19 and samples from quarter 3 were used as replacements. This change is considered to have had minimal impact on the quality of the assessment because discard rates have been consistently low in recent years.

It is not possible to reliably disaggregate the landings (and catch) data for the two subareas, and so, typically harvest rates are combined over the whole FU (Figure 2). In 2020, sampling was not carried out at the Sound of Jura because of the COVID-19 pandemic

Given the relative stock sizes and likely magnitude of the landings from the two subareas, the combined harvest rate shown in Figure 1 is expected to be more representative of the harvest rate in the Firth of Clyde than in the Sound of Jura.

Annual underwater television (UWTV) surveys are usually carried out for both subareas. The surveys have good coverage of the muddy sediment in each subarea and provide abundance estimates of each subarea with acceptable precision. The time-series for the Firth of Clyde has been continuous since 1995 and for the Sound of Jura since 2009. No survey was carried out in the Sound of Jura in 2020, and therefore an interpolated abundance estimate was used to calculate a harvest rate (average of 2019 and 2021).

Although the commercial catch-at-length samples are considered representative of the combined Norway lobster fishery in the Firth of Clyde and the Sound of Jura, sampling levels remain insufficient to provide estimates of mean weights and discard rates for the Sound of Jura separately. The discard rates and mean weights used in the catch scenarios are for the two subareas combined.




Figure 2 Norway lobster functional units in subareas 6 and 7.

### Issues relevant for the advice

During 2016–2020, the EU landing obligation was applied to all catches of Norway lobster fisheries in ICES Subarea 6 with exemptions for high survival. In 2021, this stock is still under a landing obligation and there are still exemptions in place. Observations from the 2018–2020 fishery indicate that some discarding above the minimum conservation reference size (MCRS) continues (Figure 3). Consequently, ICES is providing advice for 2022 assuming average discard rates as observed over the last three years. This is considered to be the most realistic assumption. In a situation where all catch is landed, there would be no surviving discards and the total catch advice and MSY harvest rate would be lower than those given in the catch scenario table (Table 2). However, reducing the catch of smaller Norway lobster would allow an increase in landings above those given in the catch scenario table.

The distinct areas Firth of Clyde and Sound of Jura in the FU are separated by a large area of sandy gravelly sediment around the Mull of Kintyre. These are treated as separate populations because of differences in burrow density and biological parameters, which imply different reference points.

Norway lobster in the Firth of Clyde occur at a very high density (with an average of around 0.8 individuals  $m^{-2}$ ), suggesting a relatively high productivity. The fishery in the Clyde area has been in existence since the 1960s, with the population and biological parameters having been studied numerous times. Historical harvest rates in this FU have been generally high, at or above  $F_{max}$ .  $F_{max}$  is considered an appropriate  $F_{MSY}$  proxy, expected to deliver high long-term yield with a low probability of recruitment overfishing in the Firth of Clyde. For the Sound of Jura the density is also relatively high. However, the fishery here has been sporadic and sampling is at a relatively low level; therefore, a more cautious  $F_{35\%SPR}$  (the fishing mortality that gives 35% spawning potential ratio) is considered an appropriate  $F_{MSY}$  proxy.

A single TAC covers the entire ICES Subarea 6. Management should be implemented at the FU level to ensure that fishing opportunities are in line with the scale of the resource for each of the stocks and consistent with an MSY approach.

### Reference points

**Table 4** Norway lobster in Division 6.a, Functional Unit 13. Reference points, values, and their technical basis.

#### Firth of Clyde

| Framework                    | Reference point       | Value       | Technical basis  | Source      |
|------------------------------|-----------------------|-------------|--|-------------|
| MSY approach                 | MSY $B_{trigger}$     | 580         | Lowest observed abundance estimate (Firth of Clyde); individuals in millions   | ICES (2016) |
|                              | $F_{MSY}$             | 15.1        | Proxy, harvest rate equivalent to $F_{max}$ for combined sexes, derived from a length-based per recruit analysis; percentage by number                         | ICES (2016) |
| Precautionary approach       | $B_{lim}$             | Not defined |  |             |
|                              | $B_{pa}$              | Not defined |  |             |
|                              | $F_{lim}$             | Not defined |  |             |
|                              | $F_{pa}$              | Not defined |  |             |
| EU management plan (EU 2019) | MAP MSY $B_{trigger}$ | 580         | MSY $B_{trigger}$ ; individuals in millions  | ICES (2016) |
|                              | MAP $B_{lim}$         | Not defined |  |             |
|                              | MAP $F_{MSY}$         | 15.1        | Harvest rate equivalent to $F_{MSY}$ ; percentage by number  | ICES (2016) |
|                              | MAP range $F_{lower}$ | 9.9–15.1    | Harvest rate, consistent with ranges provided by ICES, resulting in no more than 5% reduction in long-term yield compared with MSY; percentage in numbers      | ICES (2016) |
|                              | MAP range $F_{upper}$ | 15.1–15.1   | Harvest rate, $F_{MSY upper}$ value capped at $F_{MSY}$ because it has not been possible to evaluate the probability of $SSB < B_{lim}$ ; percentage by number | ICES (2016) |

**Sound of Jura**

| Framework                    | Reference point       | Value       | Technical basis  | Source      |
|------------------------------|-----------------------|-------------|--|-------------|
| MSY approach                 | MSY $B_{trigger}$     | 160         | Lowest observed abundance estimate (Sound of Jura); individuals in millions  | ICES (2016) |
|                              | $F_{MSY}$             | 12.0        | Proxy harvest rate equivalent to $F_{max}$ for combined sexes, derived from a length-based per recruit analysis; percentage by number                          | ICES (2016) |
| Precautionary approach       | $B_{lim}$             | Not defined |  |             |
|                              | $B_{pa}$              | Not defined |  |             |
|                              | $F_{lim}$             | Not defined |  |             |
|                              | $F_{pa}$              | Not defined |  |             |
| EU Management plan (EU 2019) | MAP MSY $B_{trigger}$ | 160         | MSY $B_{trigger}$ ; individuals in millions  | ICES (2016) |
|                              | MAP $B_{lim}$         | Not defined |  |             |
|                              | MAP $F_{MSY}$         | 12.0        | Harvest rate equivalent to $F_{MSY}$ ; percentage by number  | ICES (2016) |
|                              | MAP range $F_{lower}$ | 9.4–12.0    | Harvest rate, consistent with ranges provided by ICES, resulting in no more than 5% reduction in long-term yield compared with MSY; percentage in numbers      | ICES (2016) |
|                              | MAP range $F_{upper}$ | 12.0–12.0   | Harvest rate, $F_{MSY upper}$ value capped at $F_{MSY}$ because it has not been possible to evaluate the probability of $SSB < B_{lim}$ ; percentage by number | ICES (2016) |

**Basis of the assessment**

**Table 5** Norway lobster in Division 6.a, Functional Unit 13. Basis of the assessment and advice.

|                          |  |
|--------------------------|--|
| ICES stock data category | 1 (ICES, 2021b)  |
| Assessment type          | Underwater TV survey (ICES, 2021a)   |
| Input data               | One survey index (UWTV-FU13 [U6028]); commercial catches (international landings, length frequencies from Scottish and Northern Ireland catch sampling); fixed maturity parameters (from survey data); fixed natural mortalities. Discard survival rate. |
| Discards and bycatch     | Included in the assessment since 1990; dataserries from the majority of the main fleets cover almost all landings  |
| Indicators               | Size structure, mean size, and sex ratio of catches  |
| Other information        | The latest benchmark (based on the UWTV survey) was performed in 2009 (ICES, 2009)   |
| Working group            | Working Group for the Celtic Seas Ecoregion ( <a href="#">WGCSE</a> )  |

**History of the advice, catch, and management**

**Table 6** Norway lobster in Division 6.a, Functional Unit 13. ICES advice, landings, and discards. All weights are in tonnes.

| Year | ICES advice             | Landings advice for the Firth of Clyde (FU 13) | Landings advice for the Sound of Jura (FU 13) | Catch advice for the Firth of Clyde (FU 13)** | Catch advice for the Sound of Jura (FU 13)** | ICES landings | Total discards* |
|------|-------------------------|--|---|---|--|---------------|-----------------|
| 1989 |                         |  |   |   |  | 2812          |                 |
| 1990 |                         |  |   |   |  | 2909          | 193             |
| 1991 |                         |  |   |   |  | 3038          | 247             |
| 1992 | Maintain current effort |  |   |   |  | 2803          | 100             |
| 1993 | Maintain current effort |  |   |   |  | 3343          | 295             |
| 1994 | Maintain current effort |  |   |   |  | 2630          | 397             |
| 1995 | Maintain current effort |  |   |   |  | 3987          | 619             |
| 1996 | Maintain current effort |  |   |   |  | 4057          | 635             |
| 1997 | As for 1996             |  |   |   |  | 3621          | 598             |
| 1998 | Maintain current effort |  |   |   |  | 4841          | 1292            |
| 1999 | As for 1998             |  |   |   |  | 3752          | 566             |
| 2000 | Maintain current effort |  |   |   |  | 3417          | 470             |
| 2001 | As for 2000             |  |   |   |  | 3182          | 677             |
| 2002 | Maintain current effort |  |   |   |  | 3384          | 406             |
| 2003 | As for 2002             |  |   |   |  | 3173          | 1247            |
| 2004 | Maintain current effort |  |   |   |  | 2973          | 1435            |
| 2005 | As for 2004             |  |   |   |  | 3395          | 611             |

| Year | ICES advice  | Landings advice for the Firth of Clyde (FU 13) | Landings advice for the Sound of Jura (FU 13) | Catch advice for the Firth of Clyde (FU 13)** | Catch advice for the Sound of Jura (FU 13)** | ICES landings | Total discards* |
|------|--|--|---|---|--|---------------|-----------------|
| 2006 | No increase in effort  |  |   |   |  | 4780          | 515             |
| 2007 | No increase in effort and harvest rate of 15%                        | 3765   |   |   |  | 6660          | 2566            |
| 2008 | As for 2007  | 3765   |   |   |  | 5923          | 1433            |
| 2009 | No increase in effort and recent average catch                       | < 5700   |   |   |  | 4779          | 1390            |
| 2010 | Harvest rate no greater than that equivalent to fishing at $F_{0.1}$ | < 3900   |   |   |  | 5843          | 536             |
| 2011 | MSY transition scheme  | < 4100   | < 500   |   |  | 6432          | 568             |
| 2012 | MSY approach   | < 4200   | < 900   |   |  | 6687          | 1066            |
| 2013 | MSY approach   | < 5600   | < 800   |   |  | 5435          | 454             |
| 2014 | MSY approach   | < 5744   | < 521   |   |  | 6207          | 696             |
| 2015 | MSY approach   | < 3766   | < 614   |   |  | 5147          | 401             |
| 2016 | MSY approach   |  |   | ≤ 5554***                                     | ≤ 1014***                                    | 6447          | 636             |
| 2017 | MSY approach   |  |   | ≤ 5755  | ≤ 992  | 5403          | 265             |
| 2018 | MSY approach   |  |   | ≤ 4484  | ≤ 695  | 4143          | 68              |
| 2019 | MSY approach   |  |   | ≤ 5990  | ≤ 598  | 4683          | 463             |
| 2020 | Management plan  |  |   | 5227 (range 3428–5227)                        | 634 (range 496–634)                          | 3636          | 177             |
| 2021 | Management plan  |  |   | 4791 (range 3142–4791)                        | 634 (range 496–634)                          |               |                 |
| 2022 | MSY approach   |  |   | ≤ 3607  | ≤ 628  |               |                 |

\* Dead + surviving discards.

\*\* Assuming recent discard rates from 2017 onwards

\*\*\* Assuming all catches are landed

## History of catch and landings

**Table 7** Norway lobster in Division 6.a, Functional Unit 13. Catch distribution by fleet in 2020 as estimated by ICES.

| Catch       |              | Landings                      |                               | Total discards |               |
|-------------|--------------|-------------------------------|-------------------------------|----------------|---------------|
| 99% dead    | 1% surviving | <i>Nephrops</i> trawl fishery | <i>Nephrops</i> creel fishery | 75% dead       | 25% surviving |
| 3813 tonnes |              | 93% trawls                    | 7% creels                     | 177 tonnes     |               |
|             |              | 3636 tonnes                   |                               |                |               |

**Table 8** Norway lobster in Division 6.a, Functional Unit 13. History of ICES estimates of landings (for Scotland by gear) and total discards. All weights are in tonnes.

| Year | UK Scotland           |             |       |          | Other UK | Total landings | Discards* |
|------|-----------------------|-------------|-------|----------|----------|----------------|-----------|
|      | <i>Nephrops</i> trawl | Other trawl | Creel | Subtotal |          |                |           |
| 1981 | 2498                  | 404         | 66    | 2968     | 0        | 2968           |           |
| 1982 | 2372                  | 169         | 79    | 2620     | 0        | 2620           |           |
| 1983 | 3889                  | 121         | 52    | 4062     | 14       | 4076           |           |
| 1984 | 3070                  | 153         | 77    | 3300     | 10       | 3310           |           |
| 1985 | 3921                  | 293         | 65    | 4279     | 7        | 4286           |           |
| 1986 | 4073                  | 176         | 79    | 4328     | 13       | 4341           |           |
| 1987 | 2860                  | 82          | 64    | 3006     | 3        | 3009           |           |
| 1988 | 3507                  | 107         | 43    | 3657     | 7        | 3664           |           |
| 1989 | 2577                  | 184         | 35    | 2796     | 16       | 2812           |           |
| 1990 | 2731                  | 121         | 23    | 2875     | 34       | 2909           | 193       |
| 1991 | 2844                  | 145         | 26    | 3015     | 23       | 3038           | 247       |
| 1992 | 2530                  | 247         | 9     | 2786     | 17       | 2803           | 100       |
| 1993 | 3200                  | 110         | 5     | 3315     | 28       | 3343           | 295       |
| 1994 | 2503                  | 50          | 28    | 2581     | 49       | 2630           | 397       |
| 1995 | 3766                  | 131         | 26    | 3923     | 64       | 3987           | 619       |
| 1996 | 3880                  | 108         | 27    | 4015     | 42       | 4057           | 635       |
| 1997 | 3486                  | 46          | 26    | 3558     | 63       | 3621           | 598       |

| Year   | UK Scotland           |             |       |          | Other UK | Total landings | Discards* |
|--------|-----------------------|-------------|-------|----------|----------|----------------|-----------|
|        | <i>Nephrops</i> trawl | Other trawl | Creel | Subtotal |          |                |           |
| 1998   | 4540                  | 79          | 39    | 4658     | 183      | 4841           | 1292      |
| 1999   | 3476                  | 29          | 37    | 3542     | 210      | 3752           | 566       |
| 2000   | 3142                  | 63          | 75    | 3280     | 137      | 3417           | 470       |
| 2001   | 2890                  | 65          | 95    | 3050     | 132      | 3182           | 677       |
| 2002   | 3075                  | 53          | 105   | 3233     | 151      | 3384           | 406       |
| 2003   | 2954                  | 20          | 119   | 3093     | 80       | 3173           | 1247      |
| 2004   | 2619                  | 8           | 88    | 2715     | 258      | 2973           | 1435      |
| 2005   | 3148                  | 5           | 94    | 3247     | 148      | 3395           | 611       |
| 2006   | 4356                  | 1           | 179   | 4536     | 244      | 4780           | 515       |
| 2007   | 6069                  | 4           | 221   | 6294     | 366      | 6660           | 2566      |
| 2008   | 5320                  | 3           | 184   | 5507     | 416      | 5923           | 1433      |
| 2009   | 4304                  | 1           | 191   | 4496     | 283      | 4779           | 1390      |
| 2010   | 5162                  | 5           | 211   | 5378     | 465      | 5843           | 536       |
| 2011   | 5664                  | 9           | 219   | 5892     | 540      | 6432           | 568       |
| 2012   | 5617                  | 4           | 203   | 5824     | 863      | 6687           | 1066      |
| 2013   | 4708                  | 4           | 212   | 4924     | 511      | 5435           | 454       |
| 2014   | 4770                  | 1           | 258   | 5029     | 1178     | 6207           | 696       |
| 2015   | 4035                  | 8           | 206   | 4249     | 898      | 5147           | 401       |
| 2016   | 4922                  | 6           | 267   | 5195     | 1248     | 6447           | 636       |
| 2017   | 4195                  | 3           | 263   | 4461     | 942      | 5403           | 265       |
| 2018   | 3574                  | 13          | 253   | 3840     | 303      | 4143           | 68        |
| 2019** | 3834                  | 3           | 265   | 4102     | 581      | 4683           | 435       |
| 2020** | 2869                  | 10          | 225   | 3104     | 532      | 3636           | 177       |

\*Dead + surviving discards.

\*\*Landing values are preliminary

### Summary of the assessment

**Table 9** Norway lobster in Division 6.a, Functional Unit 13. Assessment summary.

| Year | UWTV abundance estimate (Clyde) | ± 2 * Standard deviations | UWTV abundance estimate (Jura) | ± 2 * Standard deviations | Landings in number | Total discards in number * | Removals in number | Harvest rate (by number)** | Landings | Total discards * | Discard proportion (by number) | Dead discard proportion (by number) | Mean weight in landings | Mean weight in discards |
|------|---------------------------------|---------------------------|--------------------------------|---------------------------|--------------------|----------------------------|--------------------|----------------------------|----------|------------------|--------------------------------|-------------------------------------|-------------------------|-------------------------|
|      | millions                        |                           |                                |                           |                    |                            |                    | %                          | tonnes   | %                |                                | grammes                             |                         |                         |
| 1995 | 579                             | 176                       | 160                            | 58                        | 207                | 82                         | 269                | 36                         | 3987     | 619              | 28                             | 23                                  | 19.24                   | 7.54                    |
| 1996 | 935                             | 242                       | 171                            | 26                        | 187                | 61                         | 233                | 21                         | 4057     | 635              | 25                             | 19.7                                | 21.68                   | 10.35                   |
| 1997 | 1198                            | 262                       | -                              | -                         | 150                | 70                         | 202                | -                          | 3621     | 598              | 32                             | 26                                  | 24.21                   | 8.50                    |
| 1998 | 1262                            | 213                       | -                              | -                         | 269                | 187                        | 409                | -                          | 4841     | 1292             | 41                             | 34                                  | 17.98                   | 6.92                    |
| 1999 | 930                             | 289                       | -                              | -                         | 216                | 93                         | 286                | -                          | 3752     | 566              | 30                             | 25                                  | 17.39                   | 6.05                    |
| 2000 | 1411                            | 246                       | -                              | -                         | 171                | 48                         | 207                | -                          | 3417     | 470              | 22                             | 17.4                                | 19.96                   | 9.75                    |
| 2001 | 1486                            | 268                       | 272                            | 76                        | 164                | 82                         | 225                | 12.8                       | 3182     | 677              | 34                             | 27                                  | 19.46                   | 8.23                    |
| 2002 | 1571                            | 288                       | 398                            | 167                       | 207                | 50                         | 245                | 12.4                       | 3384     | 406              | 19.5                           | 15.4                                | 16.35                   | 8.12                    |
| 2003 | 1817                            | 292                       | 260                            | 68                        | 166                | 134                        | 266                | 12.8                       | 3173     | 1247             | 45                             | 38                                  | 19.13                   | 9.31                    |
| 2004 | 1970                            | 367                       | -                              | -                         | 158                | 168                        | 284                | -                          | 2973     | 1435             | 52                             | 44                                  | 18.80                   | 8.54                    |
| 2005 | 1959                            | 287                       | 303                            | 84                        | 189                | 69                         | 241                | 10.7                       | 3395     | 611              | 27                             | 22                                  | 17.96                   | 8.81                    |
| 2006 | 1851                            | 257                       | 430                            | 134                       | 248                | 55                         | 290                | 12.7                       | 4780     | 515              | 18.2                           | 14.3                                | 19.27                   | 9.31                    |
| 2007 | 1233                            | 218                       | 255                            | 58                        | 350                | 387                        | 640                | 43                         | 6660     | 2566             | 53                             | 45                                  | 19.05                   | 6.64                    |
| 2008 | 1769                            | 291                       | -                              | -                         | 357                | 207                        | 512                | -                          | 5923     | 1433             | 37                             | 30                                  | 16.59                   | 6.94                    |
| 2009 | 1499                            | 210                       | 251                            | 68                        | 261                | 169                        | 388                | 22                         | 4779     | 1390             | 39                             | 33                                  | 18.31                   | 8.23                    |
| 2010 | 1750                            | 327                       | 376                            | 39                        | 276                | 55                         | 317                | 14.9                       | 5843     | 536              | 16.7                           | 13.1                                | 21.21                   | 9.68                    |
| 2011 | 2165                            | 305                       | 312                            | 73                        | 333                | 74                         | 388                | 15.7                       | 6432     | 568              | 18.2                           | 14.3                                | 19.34                   | 7.65                    |
| 2012 | 1421                            | 227                       | 371                            | 61                        | 306                | 93                         | 376                | 21                         | 6687     | 1066             | 23                             | 18.6                                | 21.83                   | 11.42                   |
| 2013 | 1990                            | 246                       | 198                            | 35                        | 262                | 62                         | 309                | 14.1                       | 5435     | 454              | 19.0                           | 15.0                                | 20.72                   | 7.37                    |




|      |      |     |     |     |     |    |     |         |      |     |      |      |       |      |
|------|------|-----|-----|-----|-----|----|-----|---------|------|-----|------|------|-------|------|
| 2014 | 1328 | 237 | 231 | 90  | 295 | 78 | 353 | 23      | 6207 | 696 | 21   | 16.6 | 20.79 | 8.92 |
| 2015 | 1820 | 351 | 376 | 127 | 232 | 54 | 273 | 12.4    | 5147 | 401 | 18.9 | 14.8 | 22.21 | 7.43 |
| 2016 | 1946 | 249 | 422 | 42  | 364 | 69 | 416 | 17.6    | 6447 | 636 | 15.9 | 12.4 | 17.70 | 9.21 |
| 2017 | 1568 | 239 | 306 | 71  | 316 | 32 | 340 | 18.1    | 5403 | 275 | 9.5  | 7.1  | 17.02 | 8.55 |
| 2018 | 2193 | 297 | 275 | 53  | 268 | 7  | 273 | 11.1    | 4143 | 68  | 2.5  | 1.90 | 16.14 | 9.79 |
| 2019 | 2083 | 381 | 318 | 61  | 271 | 64 | 319 | 13.3    | 4683 | 435 | 19.1 | 15.0 | 17.26 | 6.82 |
| 2020 | 1941 | 297 | -   | -   | 195 | 23 | 212 | 9.4 *** | 3636 | 177 | 10.7 | 8.3  | 18.96 | 7.57 |
| 2021 | 1414 | 211 | 310 | 98  | -   | -  | -   | -       | -    | -   | -    | -    | -     | -    |

\* Values prior to 2006 may be underestimates because of underreporting of landings.

\*\* Dead + surviving discards.

\*\*\* The harvest rate is estimated based on a linear interpolation of abundance for Sound of Jura in 2020, as no survey was carried out in that year.



**Figure 3** Norway lobster in Division 6.a, Functional Unit 13. The dashed lines represent catches while the solid lines represent landings. Annual length–frequency distributions are shown on the horizontal, the vertical bold lines represent mean lengths. Minimum conservation reference size (20 mm) and 35 mm visual reference levels indicated. All lengths are shown in carapace length (mm).

## Sources and references

EU. 2019. Regulation (EU) 2019/472 of the European Parliament and of the Council of 19 March 2019 establishing a multiannual plan for stocks fished in the Western Waters and adjacent waters, and for fisheries exploiting those stocks, amending Regulations (EU) 2016/1139 and (EU) 2018/973, and repealing Council Regulations (EC) No 811/2004, (EC) No 2166/2005, (EC) No 388/2006, (EC) No 509/2007 and (EC) No 1300/2008. Official Journal of the European Union, L 83: 1–17. <http://data.europa.eu/eli/reg/2019/472/oj>.

ICES. 2009. Report of the Benchmark Workshop on Nephrops (WKNEPH), 2–6 March 2009, Aberdeen, UK. ICES CM 2009/ACOM:33. 156 pp. <https://doi.org/10.17895/ices.pub.4503>.

ICES. 2016. EU request to ICES to provide  $F_{MSY}$  ranges for selected stocks in ICES subareas 5 to 10. In Report of the ICES Advisory Committee, 2016. ICES Advice 2016, Book 5, Section 5.4.1. 13 pp. <https://doi.org/10.17895/ices.advice.5612>.

ICES. 2021a. Working Group for the Celtic Seas Ecoregion (WGCSE). Draft report. ICES Scientific Reports, 3:56. 1082 pp. <http://doi.org/10.17895/ices.pub.8139>. Publication of the full report is expected at the end of October 2021

ICES. 2021b. Advice on fishing opportunities. In Report of the ICES Advisory Committee, 2021. ICES Advice 2021, section 1.1.1. <https://doi.org/10.17895/ices.advice.7720>.

[Download the stock assessment data and figures.](#)

Recommended citation: ICES. 2021. Norway lobster (*Nephrops norvegicus*) in Division 6.a, Functional Unit 13 (West of Scotland, the Firth of Clyde and Sound of Jura). In Report of the ICES Advisory Committee, 2021. ICES Advice 2021, nep.fu.13, <https://doi.org/10.17895/ices.advice.7796>.