

FICHES D'IDENTIFICATION DU ZOOPLANCTON

Editées par
J. H. FRASER

Marine Laboratory, P. O. Box 101, Victoria Road,
Aberdeen AB9 8DB, Scotland

FICHE NO. 153

(Replacing sheet No. 30)

HYDROMEDUSAE: FAMILIES ZANCLEIDAE, CLADONEMIDAE AND ELEUTHERIIDAE

by




F. S. Russell

Marine Biological Laboratory, Citadel Hill
Plymouth, Devon PL1 2 PB, England

(This sheet may be referred to in the following form:
Russell, F. S. 1977. Hydromedusae: Families Zancleidae, Cladonemidae and Eleutheriidae,
Fich. Ident. Zooplancton 153: 4 pp.)

<https://doi.org/10.17895/ices.pub.5141>

Conseil International pour l'Exploration de la Mer
Charlottenlund Slot, DK-2920 Charlottenlund
Danemark



1. *Zanclea costata*; 1a. Cnidophores on marginal tentacle. 2. *Cladonema radiatum*. 3. *Eleutheria dichotoma*.

Family Zancleidae

Mouth circular, simple. Gonads interradial. Marginal tentacles with abaxial stalked capsules, or cnidophores.

Genus *Zanclea* Gegenbaur

With two or four marginal tentacles.

1. *Zanclea costata* Gegenbaur. The medusa can be identified immediately by the characteristic structure of the marginal tentacles which carry the peculiar stalked cnidophores. The species may occur with two or four marginal tentacles.

Family Cladonemidae

Mouth with short lips armed with nematocyst clusters; stomach with periradial pouches; number of radial canals variable, some branched. Gonads completely surrounding stomach. Marginal tentacles variable in number, each with organs of adhesion.

Genus *Cladonema* Dujardin

Characters as family.

2. *Cladonema radiatum* Dujardin. Usually eight, sometimes ten, radial canals and marginal tentacles. One to four, usually three, organs of adhesion on base of each marginal tentacle. This medusa can creep or swim.

Family Eleutheriidae

Mouth circular, simple; number of radial canals variable. Gonads in special dorsal brood pouch. Marginal tentacles variable in number, each with organ of adhesion.

Genus *Eleutheria* Quatrefages

Characters as family.

3. *Eleutheria dichotoma* Quatrefages. Usually six radial canals. Usually six, deeply bifurcated marginal tentacles. Medusa buds borne on exumbrella. An inshore, creeping medusa.

FURTHER INFORMATION ON IDENTIFICATION

1. *Zanclea costata*: Allman, 1871-72, pp. 224 and 290, Pl. VII (as *Gemmaria implexa*); Hartlaub, 1907, p. 116, Figs. 104, 106-111, 115 (as *Zanclea implexa*), Figs. 115, 116 (as *Z. gemmosa*); Russell & Rees, 1936, p. 107, Figs. 1-12 (as *Z. gemmosa*); Russell, 1953, p. 99; Martin & Brinckmann, 1963.
2. *Cladonema radiatum*: Hartlaub, 1907, p. 132, Figs. 123-125; Lengerich, 1922, p. 210, Fig. 1; 1923, p. 313, Figs. G-S (as *Eleutheria radiata*); Weill, 1937, p. 438; Russell, 1953, p. 105.
3. *Eleutheria dichotoma*: Hartlaub, 1907, p. 127, Figs. 119-120; Lengerich, 1923, p. 359, Figs. R¹-Z¹, D²; Mayer, 1910, p. 94, Figs. 46-48; Russell, 1953, p. 110.

DISTRIBUTION

Species	Species
Gulf of Bothnia	—
Gulf of Finland	—
Baltic proper	—
Belt Sea	2
Kattegat	3
Skagerrak	1, 2, 3
Northern North Sea	1
Southern North Sea	1, 2, 3
English Channel (eastern)	2
English Channel (western)	1, 2, 3
Bristol Channel and Irish Sea	1, 2, 3
South and west Ireland and Atlantic	1, 2, 3
Faroe-Shetland Area	—
Faroe-Iceland Area	—
Norwegian Sea	—
Barents Sea	—

REFERENCES

- ALLMAN, GEORGE JAMES, 1871-72. Monogr. Gymnoblastic or Tubularian Hydroids. (Ray Society).
- HARTLAUB, CL., 1907. Nord. Plankt. Lief. 6, XII, Craspedote Medusen, Teil 1, Lief. 1, Codoniden und Cladonemiden, p. 1.
- LENGERICH, HANNS, 1922. Zool. Anz., Bd. LIV, Nr. 9/10, I, Wiss. Mitt. 4, p. 209.
- 1923. Zool. Jb. (Anat.), Bd. 44, Heft 3, p. 311.
- MARTIN, RAINER & BRINCKMANN, ANITA, 1963. Pubbl. Staz. zool. Napoli, 33, p. 206.
- MAYER, ALFRED GOLDSBOROUGH, 1910. Medusae of the World, Vol. I.
- RUSSELL, F. S. & REES, W. J., 1936. J. mar. biol. Ass. U.K., Vol. XXI, p. 107.
- RUSSELL, F. S., 1953. The Medusae of the British Isles, p. 99.
- WEILL, ROBERT, 1937. Bull. biol. Fr. Belg., Tome LXXI, Fasc. 4, p. 438.