

Stock Annex: Cod (*Gadus morhua*) in subareas 1 and 2 (Northeast Arctic)

Stock specific documentation of standard assessment procedures used by ICES.

Stock:	Cod
Working Group:	Arctic Fisheries Working Group (AFWG)
Created:	
Authors:	AFWG
Last benchmarked:	WKBarFar - 2021
Last updated:	March 2021
Last updated by:	AFWG – Yury Kovalev (stock coordinator), Bjarte Bogstad, Natalia Yaragina

A. General

A.1. Stock definition

The North-East Arctic cod (*Gadus morhua*) is distributed in the Barents Sea and adjacent waters, mainly in waters above 0°C. The main spawning areas are along the Norwegian coast between 67°30' and 70°N. The 0-group cod drifts from the spawning grounds eastwards and northwards and during the international 0-group survey in August it is observed over wide areas in the Barents Sea.

A.2. Fishery

The fishery for North-east Arctic cod is conducted both by an international trawler fleet operating in offshore waters and by vessels using gillnets, longlines, handlines and Danish seine operating both offshore and in the coastal areas. 60-80% of the annual landings are from trawlers. Catch quotas were introduced in the trawl fishery in 1978 and for the fisheries with conventional gears in 1989. In addition to quotas the fisheries are regulated by mesh size limitations including sorting grids, a minimum catching size, a maximum bycatch of undersized fish, maximum bycatch of non-target species, closure of areas with high densities of juveniles and by seasonal and area restrictions. Since January 1997 sorting grids have been mandatory for the trawl fisheries in most of the Barents Sea and Svalbard area. Discarding is prohibited. The minimum catching size of cod is 44 cm, and the maximum proportion of undersized fish allowed is 15% by number for cod, haddock and saithe combined. The fisheries are controlled by inspections at sea, requirement of reporting to catch control points when entering and leaving the EEZs and by inspections when landing the fish for all fishing vessels. Keeping a detailed fishing logbook on board is mandatory for most vessels, and large parts of the fleet report to the authorities on a daily basis. There is some evidence that the present catch control and reporting systems are not sufficient to prevent discarding and underreporting of catches, but it has improved considerably following the introduction of port state control in 2007.

A.3. Ecosystem aspects

Considerable effort has been devoted to investigate multispecies interactions in the Northeast Arctic. Some of these investigations have reached the stage where quantitative results are available for use in assessments. Growth of cod depends on availability of prey such as capelin (*Mallotus villosus*), and variability of cod growth has had major impacts on the cod fishery. Cod are able to compensate only partially for low capelin abundance, by switching to other prey species. This may lead to periods of high cannibalism on young cod, and may result in impacts on other prey species which are greater than those estimated for periods when capelin is abundant. In a situation with low capelin abundance, juvenile herring (*Clupea harengus*) experience increased predation mortality by cod. The timing of cod spawning migrations is influenced by the presence of spawning herring in the relevant area. The interaction between capelin and herring is illustrated by the recruitment failure of capelin coinciding with years of high abundance of young herring in the Barents Sea. Herring predation on capelin larvae is believed to be partially responsible for the recruitment failure of capelin when young herring are abundant in the Barents Sea.

The composition and distribution of species in the Barents Sea depend considerably on the position of the polar front which separates warm and salty Atlantic waters from colder and fresher waters of arctic origin. Variation in the recruitment of some species including cod and capelin has been associated with the changes in the influx of Atlantic waters to the large areas of the Barents Sea shelf.

The annual consumption of herring, capelin and cod by marine mammals (mainly harp seals and minke whales) has been estimated to be in the order of 1.5–2.0 million tonnes (Bogstad *et al.*, 2000; See also Table 1.9 ICES 2014).

However, estimates of total annual food consumption of Barents Sea harp seals are in the range of about 3.3–5 million tonnes (depending on choice of input parameters, Nilssen *et al.*, 2000). The applied model used different values for the field metabolic rate of the seals (corresponding to two or three times their predicted basal metabolic rate) and under two scenarios: with an abundant capelin stock and with a very low capelin stock.

If capelin was abundant the total harp seal consumption was estimated to be about 3.3 million tons (using lowest field metabolic rate). The estimated consumption of various commercially important species was as follows (in tonnes): capelin approximately 800 000, polar cod (*Boreogadus saida*) 600 000, herring 200 000 and Atlantic cod 100 000.

A low capelin stock in the Barents Sea (as it was in 1993–1996) led to switches in seal diet composition, with estimated increased consumption of polar cod (870 000 tonnes), other gadoids (mainly Atlantic cod; 360 000 tonnes), and herring (390 000 tonnes).

B. Data

B.1. Commercial catch

Norway

Norwegian commercial catch in tonnes by quarter, area and gear are derived from the sales notes statistics of The Directorate of Fisheries. Data from about 20 subareas are aggregated on 6 main areas for the gears gillnet, longline, handline, purse-seine, Danish seine, bottom trawl, shrimp trawl and trap. For bottom trawl the quarterly area distribution of the catches is adjusted by logbook data from The Directorate of Fisheries

and the total bottom-trawl catch by quarter and area is adjusted so that the total annual catch for all gears is the same as the official total catch reported to ICES.

No discards are reported or accounted for, but there are several reports of discards.

The sampling strategy is to have age and length samples from all major gears in each main area and quarter. The main sampling program is sampling the landings. Additional samples from catches are obtained from the IMR reference fleet (fishing vessels contracted for sampling), and the coast guard.

The ECA software (Hirst *et al.*, 2012) has been developed to utilize all sampling information to estimate catch-at-age for areas (1, 2a and 2b), quarters and gears (bottom trawl, gillnet, Danish seine and longline/handline). This software also handles the splitting of catches into NEA cod and Coastal cod. A revision of the Norwegian catch-at-age series is planned for 2022, preliminary information about this revision can be found in WD6, WKBarFar 2021).

Russia

Russian commercial catch in tonnes by quarter and area are derived from the All-Russian Institute of fishery and oceanography (Moscow) statistics department. Data from each fishing vessel are aggregated on three ICES subdivisions (1, 2a and 2b). Russian fishery by passive gears was almost stopped by the end of the 1940s. At present the bottom-trawl fishery constitutes more than 95% of the cod catch.

The sampling strategy was to conduct length and weight measurements and collect age samples directly at sea, on board of both research and commercial vessels to have age and length distributions from each area and quarter. Data on length distribution of cod in catches were collected in areas of cod fishery all year round by a "standard" fishery trawl (since 2011 the mesh size is 130 mm for the entire Barents Sea, previously it was 125 mm in the Russian Economic zone and Svalbard area and 135 mm in the Norwegian Economic zone) and summarized by three ICES Subareas (1, 2a and 2b). Previously the PINRO area divisions were used, which differed from the ICES subdivisions.

Age sampling was carried out by two ways: without any selection (otoliths were taken from any fish caught in one trawl haul, usually from 100–300 sp.) or using a stratified by length sampling method (i.e. approximately 10–15 sp. per each 10-cm length group). The last method has been used since 1988.

All fish taken for age-reading were measured and weighted individually.

Catch-at-age are reported to ICES AFWG by subdivision (1, 2.a and 2.b) and quarter (before 1984 – by subdivision and year). Data on length distribution of cod in catches, as well as age-length keys, are formed for each quarter and area. In the case when a catch is present in the area/quarter but a length frequency is absent, a length frequency for the corresponding quarter, summarized for the whole sea is used. If there is no data on length composition of cod in catches per a quarter within the whole sea, a frequency summarized for the whole year and whole sea is used. Gaps in age-length distributions in subdivisions are filled in with data from the corresponding quarter, summarized for the whole sea. Remaining gaps are filled in with information from the age-length key formed for the long-term period (1982–2018) for each quarter and for the whole sea (Kovalev and Yaragina, 1999; WD 11 WKBarFar, 2021). Before 1982, calculation of catch-in-numbers in subdivisions was based on the age-length keys for the whole year and length distribution in catches.

Germany and Spain

Catch-at-age is reported to the WG by ICES subdivision (1, 2.a and 2.b) and quarter, according to national sampling. Missing quarters/subdivisions are filled in by use of Russian or Norwegian sampling data.

Other nations

Total annual catch in tonnes is reported by ICES subdivisions. All catches by other nations are taken by trawl. The age composition from the sampled trawl fleets is therefore applied to the catches by other nations. Reported catch in tonnes to ICES by other nations is supplemented and sometimes adjusted by reported catches to Norwegian and Russian authorities. Time series of catches by Faroes and Greenland are under revision (WD6, WKBarFar 2021).

The text table below shows which country supplied which kind of data for 2019:

Country	KIND OF DATA				
	Caton (catch in weight)	Canum (catch-at-age in numbers)	Weca (weight at age in the catch)	Matprop (proportion mature by age)	Length composition in catch
Norway	x	x	x	x	x
Russia	x	x	x	x	x
Germany	x	x	x		
UK	x				
France	x				
Spain	x	x	x		x
Portugal	x				
Poland	x	x	x		x
Greenland	x				
Faroe Islands	x				
Iceland	x				
Belarus	x				
Lithuania	x				
Latvia	x				
Estonia	x				

Since 2008 the catch data has been handled by Intercatch. Earlier the nations that sample the catches, provided the catch-at-age data and mean weights at age on Excel spreadsheet files, and the national catches were combined in Excel spreadsheet files. Historic data should be found in the national laboratories and with the stock coordinator. Data (excel files) back to 1993 are available on WkbarFar 2021 Sharepoint and collection and computerisation of historic data farther back in time is ongoing.

For 1983 and later years mean weight at age in the catch is calculated as the weighted average for the sampled catches for age 1-11.

For age 12 and older, the mean weight at age data was reconstructed using the von Bertalanffy growth equation for both data series weight in stock and weight in catch. The method is presented in WD 12 (WKBarFar 2021). Mean weights at age are fitted by cohorts (starting from the year class 1981) using WECA data for ages 3-11 years; parameters t_0 and W_∞ of the von Bertalanffy equation are fitted as independent of time (to all generations except some outliers). Some year classes (YC) with the most noisy data were excluded from the fitting of t_0 and W_∞ (e.g. 1981 1989, 1992, 1995).

For the earlier period (1946–1982) mean weight at age in catches is set equal to mean weight at age in the stock (ICES 2001). Full data set including new WECA data for ages 12-15+ are presented in AFWG 2021 Table 3.8.

B.2. Biological

Weight in stock and maturity

For 1983 and later years weight at age in the stock and maturity-at-age is calculated as weighted averages from Russian and Norwegian surveys during the winter season. Stock weights at age a (W_a) at the start of year y for ages 1-11 are calculated as follows:

$$W_{a,y} = 0.5(W_{rus,a-1,y-1} + (\frac{N_{nbar,a,y}W_{nbar,a,y} + N_{lof,a,y}W_{lof,a,y}}{N_{nbar,a,y} + N_{lof,a,y}})) \tag{1}$$

where

$W_{rus,a-1}$: Weight at age $a-1$ in the Russian survey in year $y-1$

$N_{nbar,a}$: Abundance at age a in the Norwegian Barents Sea acoustic survey in year y

$W_{nbar,a}$: Weight at age a in the Norwegian Barents Sea trawl survey in year y

$N_{lof,a}$: Abundance at age a in the Lofoten survey in year y

$W_{lof,a}$: Weight at age a in the Lofoten survey in year y

When the Russian survey was discontinued in 2018 (it was not conducted in 2016 either) it was agreed to use mean long term correction factors that would allow using only the weight-at-age from the Joint Barents Sea and Lofoten surveys. The correction factors to calculate weight and maturity at age were updated during the 2019 AFWG meeting (see Working document No.15), based on historical differences between Winter survey+Lofoten and Russian survey data, following the approach used in 2017 (Yaragina and Bogstad, AFWG-2017, WD 10).

Correction factors for stock weights at age are as follows (WD 15 AFWG 2019):

AGE, YEARS	1	2	3	4	5	6	7	8	9	10	11
Mean Ratio between the combined data and the Joint Barents Sea + Lofoten surveys data (calculated based on 1985-2016, 2018 data)	0.93	0.93	0.88	0.88	0.93	0.95	0.96	0.98	1.01	1.01	1.10

At the WKBarFar meeting (February 2021), it was decided to use also the Ecosystem survey (BESS) data (since 2004) on mean weight at age (ages 2-10 years) in order to reduce random fluctuations that might increase because of discontinuation of Russian Autumn survey.

The following approach is used for ages 3-11 (WD 12, WKBarFar 2021):

$$New_west_{a,y} = SW_{sur} * WEST_{a,y} * + SW_{sur} * EcoWeight_{a-1,y-1} * mean_corr_factor_a \tag{2}$$

where SW_{sur} is weighting coefficient for each data source,

$WEST_{a,y}$ is mean weight at age in stock calculated by equation 1,

$Eco_Weight_{a-1,y-1}$ is mean weight at age from Ecosystem survey taken for the same cohort,

$mean_corr_factor_a$ is a coefficient of proportionality between $WEST_{a,y}$ and Weight-at-age $_{a-1,y-1}$ from Ecosystem survey, representing mean growth in weight from the time of Eco survey (August –September) and assessment time (1 January of the next year).

Equal weight should be given to each survey. Thus, the SW_{sur} assumed for each survey = 0.33 if we are combining three surveys (means =0.67 for WEST data in cases where they include Russian Autumn survey) and 0.5, if we combine two (in cases where WEST does not include Russian Autumn survey). The Eco survey data will have a SW_{sur} = 0.33 in the first case and 0.5 in the second one.

Russian Autumn survey has not been conducted since 2018 (and in 2016 also); data of the BESS from 2014 and 2018 were not used because of incomplete survey coverage.

Correction factors for the BESS stock weights at age accounting for mean growth from August-September to 1st January of the next year are as follows (WD 12 WKBarFar 2021), shown in the text table below: These factors should be updated at benchmark meetings.

MEAN RATIOS (CORRECTION FACTORS) BETWEEN THE COMBINED DATA ON WEIGHT AT AGE (FROM EQUATION 1) AND DATA FROM ECOSYSTEM SURVEY CALCULATED BY STOX:AGE, YEARS	3	4	5	6	7	8	9	10	11
Mean Ratio between the combined data and the Ecosystem survey (a-1,y-1) (based on data for 2004-2019 except 2014 and 2018)	1.56	1.36	1.22	1.19	1.19	1.21	1.23	1.23	1.19

Mean weights of NEA cod in stock for ages 12 and older are now calculated via von Bertalanffy growth equation using the method presented in WD 12 (WKBarFar 2021) instead of previously used fixed values. Mean weights at age are fitted by cohorts (starting from the year class 1981) using WEST data for ages 3-11 years; parameters t_0 and W_∞ of the von Bertalanffy equation are fitted as independent on time (to all generations except some outliers). Some year classes with the most noisy data were excluded from the fitting t_0 and W_∞ (e.g. 1981-1983, 1987-1989, 1992, 1994).

New WEST data for ages 3-15+ are presented in AFWG 2021 Table 3.9.

Maturity-at-age is estimated from the same surveys by the same formulae (1), replacing weight by proportion mature.

As the Russian survey was discontinued in 2018 (it was not conducted in 2016 either) it was agreed to use mean long term correction factors that would allow using only the maturity-at-age from the Joint Barents Sea and Lofoten surveys.

Correction factors for stock maturity at age are as follows (WD 15 AFWG 2019):

AGE, YEARS	3	4	5	6	7	8	9	10	11	12
Mean Ratio between the combined data and the Joint Barents Sea + Lofoten surveys data (calculated based on 1989-2016, 2018 data)	0.88	0.94	0.82	0.83	0.91	0.96	0.98	0.99	1.00	1.00

The time series for weight and maturity at age was revised in 2021 following the revision of the time series in the Norwegian (Joint) winter survey (see WD 1 WKBarFar 2021) and Lofoten survey. For the earlier period (1946–1982) the maturity-at-age and

weight at age in the stock is based on Russian sampling in late autumn (both from fisheries and from surveys) and Norwegian sampling in the Lofoten spawning fishery. These data were introduced and described in the 2001 assessment report (ICES 2001).

Natural mortality/cannibalism

Natural mortality (M) is assumed to be equal to 0.2 plus cannibalism mortality for ages 1–6.

The method used for calculation of the prey consumption by cod described by Bogstad and Mehl (1997) is used to calculate the consumption of cod by cod for use in SAM. The consumption is calculated based on cod stomach content data taken from the joint PINRO-IMR stomach content database (methods described in Mehl and Yaragina, 1992). On average about 9000 cod stomachs from the Barents Sea have been analysed annually in the period 1984–2019.

These data are used to calculate the per capita consumption of cod by cod for each half-year (by prey age groups 3–6 and predator age groups 3-11+). It was assumed that the mature part of the cod stock is found outside the Barents Sea for three months during the first half of the year. Thus, consumption by cod in the spawning period was omitted from the calculations.

The number of cod predators at age is taken from the SAM, and thus an iterative procedure has to be applied. All occurrences of intra-cohort predation were removed from the dataset as these could possibly cause problems with convergence. The following procedure realized in R script was followed:

The 1st run of SAM is done without taking into account consumption with a mortality of 0.2. It gives a stock numbers, simulated catches and fishing mortality.

Stage 2 - on the basis of the equation (1) which is similar to Pope's approximation, we calculate the number of the corresponding cohorts in terms of both consumption and catch (ages 3-6) for the period 1984- the last year and $M=0.2$:

$$N(y,a) = N(y+1,a+1) * e^M + C(y,a) * e^{M/2} + \text{Cons}(y,a) * e^{M/2} \quad (1)$$

Where: $N(y, a)$ is the number of cod at the age a , at the beginning of the year y ,

$M = 0.2$,

$C(y, a)$ - simulated catch from first run,

$\text{Cons}(y,a)$ - the amount of young cod consumed at ages 3-6 and the period 1984-the last year calculated using the abundance of older cod from the last run and data on per capita annual consumption of cod

Stage 3 - Calculate the mortality rate from cod predation according to the equation (2):

$$M2(y,a) = \ln(N(y,a) / N(y+1,a+1)) - F(y,a) - 0.2 \quad (2)$$

$F(y,a)$ - fishing mortality from last run.

Stage 4 Run the model with the natural mortality taken from the calculated mortality from predation ($0.2 + M2$).

Stage 5 – repeat from stage 2 onwards until difference between summary of cod abundance at ages 3-6 for period 1984-the last year from last iteration and previous ones becomes sufficiently low (formal convergence criteria was not used, in practice 5 iterations was used).

The process noise in N is captured in the SAM model. It may cause situations when the ad-hoc M_s becomes less than 0.2. In the current implementation they are then set to 0.2.

A process model for M_2 should be developed.

Since 2015 hindcasted data on cod cannibalism for the historical period (1946–1983) are also available. These have been applied to make the VPA time-series with cannibalism consistent (Yaragina *et al.*, WD7 WKARCT 2015).

Both the proportion of natural mortality before spawning (M_{prop}) and the proportion of fishing mortality before spawning (F_{prop}) are set to 0. The peak spawning in the Lofoten area occurs most years in late March-early April.

B.3. Surveys

General: Survey data should be made available separate by age up to age 15+. At present the oldest age used in the tuning is 12+ (changed from 12 at WkBarFar 2021).

Russia

Russian surveys of cod in the southern Barents Sea started in the late 1940s as trawl surveys of young demersal fish. Since 1957 such surveys have been conducted over the whole feeding area including the Bear Island - Spitsbergen area (Baranenkov, 1964; Trambachev, 1981), both young and adult cod have been surveyed simultaneously. In 1984, acoustic methods started to be implemented during surveys of fish stocks (Zaferman and Serebrov, 1984; Lepesevich, Shevelev, 1997; Lepesevich *et al.*, 1999). In 1995 a new acoustic assessment method was applied for the first time, which allowed the differentiation and registration of echo intensities from fish of different length (Shevelev *et al.*, 1998). Methods of calculations of survey indices also changed, e.g. due to the necessity to derive length-based indices for the FLEKSIBEST model (Bogstad *et al.*, 1999; Gusev and Yaragina, 2000).

Survey duration has been reduced from 5–6 months (September-February) in 1946–1981 to 2–2.5 months (October-December) since 1982. The aim of conducting a survey is to investigate both the commercial size cod as well as the young cod and to receive reliable data to compose annual maturity ogives. The survey covers the main areas where juveniles settle down as well as the commercial fishery takes place, including cod at age 0+ - 10+ years. A total of more than 400 trawl hauls are conducted during the survey (mainly bottom-trawl hauls, a few pelagic trawl hauls).

There are two survey abundance indices at age: 1) absolute numbers (in thousands) computed from the acoustics and 2) trawl swept-area indices, calculated as absolute numbers registered in survey standard area (Golovanov *et al.*, 2006, 2007). The Russian survey was discontinued in 2018.

Joint Russian-Norwegian winter (February) survey

The survey started in 1981 and covers the ice-free part of the Barents Sea. Both swept-area estimates from bottom-trawl and acoustic estimates are produced. The swept-area estimates are used in the tuning for ages 3–8, and the acoustic estimate are added to the Norwegian acoustic survey in Lofoten and used for tuning for ages 3–9. The survey is described in Jakobsen *et al.*, (1997) and Mehl *et al.*, (2013, 2014). The time series for the bottom trawl estimates has recently been revised back to 1994 (Mehl *et al.* 2016). Both acoustic and bottom trawl indices of the survey were revised (1994-2020) using bootstrap mean estimates and including three northern strata (24-26) in the indices (WD 1 WKBarFar 2021).

Norwegian Lofoten survey

Acoustic estimates from the Lofoten survey extends back to 1985. The survey is described by Korsbrekke (1997). The indices were revised for years 2010-2020 (WD 2 WKBarFar 2021).

Joint Russian–Norwegian Ecosystem survey (August–September)

This survey started in 2004, but is a continuation and integration of previous surveys conducted at this time of year (0-group survey, capelin survey, various bottom-trawl investigations). The survey methodology and results are described in annual survey reports (e.g. Prokhorova 2013). Unfortunately, there is at present no agreed method for calculating bottom-trawl indices from this survey (Dingsør, WD17, WKARCT 2015 vs. ICES AFWG 2014 Table A14). Agreeing on a common methodology has very high priority.

Commercial cpue

Norwegian and Russian commercial CPUE data are made available for AFWG each year but are not used in the assessment.

C. Estimation of historical stock development

Model used: SAM (Interbenchmark in 2017, ICES 2017, Updated by benchmark 2021, ICES 2021).

Input data types and characteristics:

TYPE	NAME	YEAR RANGE	AGE RANGE	VARIABLE FROM YEAR TO YEAR YES/NO
Caton	Catch in tonnes	1946–last data year		Yes
Canum	Catch-at-age in numbers	1946 –last data year	3 –15+	Yes
Weca	Weight at age in the commercial catch From observations Restored using von Bertalanffy growth equation	1982–last data year	3–15+	Yes, set equal to west for 1946-1981 for all ages
			3-11	
			12-15+	
West	Weight at age of the stock at spawning time. From observations Restored using von Bertalanffy growth equation	1946–last data year	3–15+	Yes
			3-11	
			12-15+	
Mprop	Proportion of natural mortality before spawning	1946–last data year	3–15+	No, set to 0 for all ages in all years
Fprop	Proportion of fishing mortality before spawning	1946–last data year	3–15+	No, set to 0 for all ages in all years

Matprop	Proportion mature at age	1946–last data year	3–15+	Yes
Natmor	Natural mortality	1946–last data year	3–15+	No, values 0.2 for all ages in all years
	Additional natural mortality caused by cannibalism	1946–last data year	3–6	Yes, annual est. of cannibalism from 1984, for period 1946-1983 set to hindcasted values since 2015 WG (WD7 WKARCT 2015)

Tuning data

TYPE	NAME	YEAR RANGE	AGE RANGE
Tuning fleet 1	Joint Barents Sea survey, February Splitted into two periods: 1981-2013 and 2014-last data year	1981–last data year	3–12+
Tuning fleet 2	Joint Barents Sea Acoustic, February+ Lofoten Acoustic survey in March	1985–last data year	3-12+
Tuning fleet 3	Russian bottom-trawl survey, November	1984–last data year	3–12+
Tuning fleet 4	Barents Sea ecosystem survey, September	2004–last data year	3–12+

SAM settings

The order of fleets in sets of parameters:

Catches

Tuning fleet 1 (first part)

Tuning fleet 1 (second part)

Tuning fleet 2

Tuning fleet 3

Tuning fleet 4

\$minAge

The minimum age class in the assessment

3

\$maxAge

The maximum age class in the assessment

15

\$maxAgePlusGroup

Is last age group considered a plus group (1 yes, or 0 no).

1 1 1 1 1 1

\$keyLogFsta

Coupling of the fishing mortality states (normally only first row is used).

0 1 2 3 4 5 6 7 8 9 10 11 11
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1

\$corFlag

Correlation of fishing mortality across ages (0 independent, 1 compound symmetry, or 2 AR(1))

0

\$keyLogFpar

Coupling of the survey catchability parameters (normally first row is not used, as that is covered by fishing mortality).

-1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1
 0 1 2 3 4 5 6 7 8 8 -1 -1 -1
 9 10 11 12 13 14 15 16 17 17 -1 -1 -1
 18 19 20 21 22 23 24 25 26 26 -1 -1 -1
 27 28 29 30 31 32 33 34 35 35 -1 -1 -1
 36 37 38 39 40 41 42 43 44 44 -1 -1 -1

\$keyQpow

Density dependent catchability power parameters (if any).

-1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1

\$keyVarF

Coupling of process variance parameters for log(F)-process (normally only first row is used)

0 1 1 1 1 1 1 1 1 1 1 1 1
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1
 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1

\$keyVarLogN

Coupling of process variance parameters for log(N)-process

0 1 1 1 1 1 1 1 1 1 1 1 1

`$keyVarObs`

Coupling of the variance parameters for the observations.

```
0 1 2 2 2 2 2 2 3 3 4 4 4
5 6 6 6 6 7 7 7 7 7 -1 -1 -1
5 6 6 6 6 7 7 7 7 7 -1 -1 -1
8 8 8 8 8 8 9 9 9 9 -1 -1 -1
10 10 10 10 10 10 11 11 11 11 -1 -1 -1
12 12 12 12 12 12 12 12 12 12 -1 -1 -1
```

`$obsCorStruct`

Covariance structure for each fleet ("ID" independent, "AR", AR(1), or "US" for unstructured). | Possible values are: "ID" "AR" "US"

```
"ID" "AR" "AR" "AR" "AR" "AR"
```

`$keyCorObs`

Coupling of correlation parameters can only be specified if the AR(1) structure is chosen above.

NA's indicate where correlation parameters can be specified (-1 where they cannot).

```
#3-4 4-5 5-6 6-7 7-8 8-9 9-10 10-11 11-12 12-13 13-14 14-15
```

```
NA NA NA NA NA NA NA NA NA NA NA NA NA
```

```
0 0 0 1 1 2 2 3 -1 -1 -1
0 0 0 1 1 2 2 3 -1 -1 -1
4 4 4 5 6 6 6 7 8 -1 -1 -1
9 9 9 9 10 10 10 11 -1 -1 -1
12 12 12 13 13 13 14 14 15 -1 -1 -1
```

`$stockRecruitmentModelCode`

Stock recruitment code (0 for plain random walk, 1 for Ricker, and 2 for Beverton-Holt).

```
0
```

`$noScaledYears`

Number of years where catch scaling is applied.

```
0
```

`$keyScaledYears`

A vector of the years where catch scaling is applied.

`$keyParScaledYA`

A matrix specifying the couplings of scale parameters (nrow = no scaled years, ncols = no ages).

`$fbarRange`

lowest and highest age included in Fbar

```
5 10
```

`$keyBiomassTreat`

To be defined only if a biomass survey is used (0 SSB index, 1 catch index, and 2 FSB index).

```
-1 -1 -1 -1 -1 -1
```

\$obsLikelihoodFlag

Option for observational likelihood | Possible values are: "LN" "ALN"
"LN" "LN" "LN" "LN" "LN" "LN"

\$fixVarToWeight

If weight attribute is supplied for observations this option sets the treatment (0 relative weight, 1 fix variance to weight).

0

Model options chosen for XSA

(used as an additional model for checking of results):

Tapered time weighting applied, power = 3 over 20 years

Catchability independent of stock size for ages > 12

Catchability independent of age for ages > 12

Survivor estimates shrunk towards the mean F of the final 5 years or the 3 oldest ages

S.E. of the mean to which the estimate are shrunk = 1.5

Shrinkage to the population mean (p-shrinkage) not applied

Minimum standard error for population estimates derived from each fleet = 0.3

Prior weighting not applied

D. Short-term projection

Model used: Age structured

Software used: R script prediction with management option table SAM

Initial stock size (intermediate year): Taken from SAM for age 4 and older. The recruitment-at-age 3 for the intermediate year and the following 2 years are estimated from survey data and environmental data using the "hybrid model" described in section 1.4.2 in AFWG 2018 (ICES 2018)

Natural mortality: average of the three last years or set equal to the values estimated for the terminal year if there is a strong trend during the most recent years.

Maturity: average of the three last years

F and M before spawning: Set to 0 for all ages in all years

Weight at age in the stock: Predicted by applying 3-year average of annual increments by cohort on last year's observation.

Weight at age in the catch: Predicted by applying 5-year average annual increments by cohort on last year's observation, for intermediate year and intermediate year+1. For intermediate year+2 and later years weight at age in catch is set equal to the values for intermediate year+1.

Exploitation pattern: Average of the recent years taking into account stability of the pattern. 5 years average as default.

Intermediate year assumptions: Normally F status quo is used. If this corresponds to a catch which deviates considerably from the agreed TAC, one should consider other approaches.

Stock recruitment model used: None

Procedures used for splitting projected catches: Not relevant

E. Medium-term projections

F. Long-term projections

MSY and HCRs in long-term perspective have been investigated during special workshops (ICES 2015, ICES 2016) using long-term stochastic simulations. The model was updated and using most recent data and simulations were redone during the Benchmark 2021 (ICES 2021). The PA and MSY reference points were considered to be adequate and the previously used values were not changed.

G. Biological reference points

Introduced 1998: $B_{lim}=112\ 000\ t$, $B_{pa}=500\ 000\ t$, $F_{lim}=0.70$, $F_{pa}=0.42$

Adopted in 2003: $B_{lim}=220\ 000\ t$, $B_{pa}=460\ 000\ t$, $F_{lim}=0.74$, $F_{pa}=0.40$

F_{MSY} is estimated to be in the range 0.4-0.6

MSY $B_{trigger}$ is at the level of 460 000 t (B_{pa}), and used as a trigger point in HCR.

H. Harvest control rule

The HCR adopted by JNRFC in 2016 following evaluation of several rules in 2016 (ICES 2016) is a two-step rule where F increases if SSB goes above $2 \times B_{pa}$. The rule now reads as follows:

The TAC is calculated as the average catch predicted for the coming 3 years using the target level of exploitation (F_{tr}).

The target level of exploitation is calculated according to the spawning stock biomass (SSB) in the first year of the forecast as follows:

- if $SSB < B_{pa}$, then $F_{tr} = SSB / B_{pa} \times F_{msy}$;
- if $B_{pa} \leq SSB \leq 2 \times B_{pa}$, then $F_{tr} = F_{msy}$;
- if $2 \times B_{pa} < SSB < 3 \times B_{pa}$, then $F_{tr} = F_{msy} \times (1 + 0.5 \times (SSB - 2 \times B_{pa}) / B_{pa})$;
- if $SSB \geq 3 \times B_{pa}$, then $F_{tr} = 1.5 \times F_{msy}$;

where $F_{msy}=0.40$ and $B_{pa}=460\ 000$ tonnes.

If the spawning stock biomass in the present year, the previous year and each of the three years of prediction is above B_{pa} , the TAC should not be changed by more than +/- 20% compared with the previous year's TAC. In this case, F_{tr} should however not be below 0.30.

H. Other issues

1. References

- Baranenkov, A.S. 1964. Some results of estimation of cod fry in the Barents Sea during 1946-1961. *In: Materials of PINRO Scientific Council Meeting by the results of investigations conducted in 1962-63. PINRO, Murmansk, 3:72-107 (in Russian).*
- Bogstad B., Fotland Å., Mehl S., 1999. A revision of the abundance indices for cod and haddock from the Norwegian winter survey in the Barents Sea, 1983-1999. Working Document N 16 for the Arctic Fisheries Working Group, August 1999, 14 pp.
- Bogstad, B., Haug, T. and Mehl, S. 2000. Who eats whom in the Barents Sea? NAMMCO Scientific Publications 2: 98-119.
- Bogstad, B. and Mehl, S. 1997. Interactions Between Cod (*Gadus morhua*) and Its Prey Species in the Barents Sea. *Forage Fishes in Marine Ecosystems. Proceedings of the International Symposium on the Role of Forage Fishes in Marine Ecosystems. Alaska Sea Grant College Program Report No. 97-01: 591-615. University of Alaska Fairbanks.*
- Dingsør, G. E. 2015. Bottom-trawl indices from ecosystem survey for cod and haddock, using a design-based approach. WD17, WKARCT 2015.
- Golovanov, S.E., Yaragina, N.A. and Sokolov, A.M. 2006. Revised indices of the Northeast Arctic cod abundance according to the data from Russian trawl-acoustic survey (TAS) for bottom fish species. WD #21 for AFWG 2006.
- Golovanov S.E., Sokolov A.M., and Yaragina N.A. 2007. Revised indices of the Northeast Arctic cod abundance according to the 1982-2006 data from Russian trawl-acoustic survey (TAS). Working Document #3 for AFWG 2007.
- Gusev E.V. and Yaragina N.A. 2000. A conversion of cod abundance from the trawl-acoustic survey data to a number of cod in different age-length groups to be used in the Flexibest model. Working Document N 35 for the Arctic Fisheries Working Group, August 2000, 9 pp.
- Hirst, D., Storvik, G., Rognebakke, H., Aldrin, M., Aanes, S., and Vølstad, J. H. 2012. A Bayesian modelling framework for the estimation of catch-at-age of commercially harvested fish species. *Can. J. Fish. Aquat. Sci.* 69: 2064–2076.
- ICES 2001. Report of the Arctic Fisheries Working Group. Bergen, Norway, 24 April – 3 May 2001. ICES CM 2001/ACFM:19. Part 2. 380 pp.
- ICES 2003. Report of the Study Group on Biological Reference Points for Northeast Arctic Cod, Svanhovd, Norway 13-17 January 2003. ICES C. M. 2003/ACFM:11, 39 pp.
- ICES. 2014. Report of the Arctic Fisheries Working Group (AFWG),, 23-29 April 2014, Lisbon, Portugal. ICES CM 2014/ACOM:05. 656 pp.
- ICES 2015. Report of the Benchmark Workshop on Arctic Stocks (WKARCT), Copenhagen 26-30 January 2015. ICES C. M. 2015/ACOM:31, 125 pp.
- ICES. 2016. Report of the second workshop on Management Plan Evaluation on North-east Arctic cod and haddock and Barents Sea capelin (WKNEAMP-2), 25–28 January 2016, Kirkenes, Norway. ICES CM 2016/ACOM:47. 76 pp.
- ICES. 2017. Report of the Inter-Benchmark Protocol on Northeast Arctic cod (IBPArcticCod), 4–6 April 2017, ICES HQ, Copenhagen, Denmark. ICES CM 2017/ACOM:29.
- ICES 2018. Report of the Arctic Fisheries Working Group, Ispra, Italy, 16-24 April 2018. ICES C.M. 2018/ACOM:06,857 pp.
- ICES. 2021. Benchmark Workshop for Barents Sea and Faroese Stocks (WKBARFAR 2021). ICES Scientific Reports. 3:21. xx pp. <https://doi.org/10.17895/ices.pub.7920>.
- Jakobsen, T., Korsbrekke, K., Mehl, S., and Nakken, O. 1997. Norwegian combined acoustic and bottom-trawl surveys for demersal fish in the Barents Sea during winter. ICES CM 1997/Y:17.

- Korsbrekke, K. 1997. Norwegian acoustic survey of Northeast Arctic cod on the spawning grounds off Lofoten. ICES C.M 1997/Y:18.
- Kovalev, Y., and Bogstad, B. 2005. Evaluation of maximum long-term yield for Northeast Arctic cod. In Shibarov, V. (ed.): Proceedings of the 11th Joint Russian-Norwegian Symposium: Ecosystem dynamics and optimal long-term harvest in the Barents Sea fisheries. Murmansk, Russia 15-17 August 2005. IMR/PINRO Report series 2/2005, p. 138-157.
- Kovalev, Yu. A. and Bogstad, B. 2015. Update of the population model for Northeast Arctic cod. WD6, WKARCT 2015.
- Kovalev Yu., Yaragina N.A., 1999. Scheme of recalculation of cod commercial catch weight into number of fish of different length-age groups accepted in PINRO (Russia) for the Flexibest model. Working Document N 10 for the Arctic Fisheries Working Group, August 1999, 9 pp.
- Lepesevich, Yu. M. and Shevelev, M. S. 1997. Evolution of the Russian survey for demersal fish: From ideal to reality. ICES C. M. 1997/Y:09.
- Lepesevich Yu.M., Smirnov O.V. and K.V. Drevetnyak, 1999. The Russian trawl acoustic survey on demersal adult and young fish stock assessments in the Barents Sea in autumn/winter. Working Document No 7 for the Arctic Fisheries Working Group, August 1999, 11 pp.
- Mehl, S., Aglen, A., and Johnsen, E. 2016. Re-estimation of swept area indices with CVs for main demersal fish species in the Barents Sea winter survey 1994 – 2016 applying the Sea2Data StoX software. Fisker og Havet 10-2016. Institute of Marine Research, Bergen, Norway.
- Mehl, S., Aglen, A., Alexandrov, D.I., Bogstad, B., Dingsør, G.E., Gjørseter, H., Johannesen, E., Korsbrekke, K., Murashko, P.A., Prozorkevich, D.V., Smirnov, O., Staby, A., and Wenneck, T. de Lange, 2013. Fish investigations in the Barents Sea winter 2007-2012. IMR-PINRO Joint Report Series 1-2013, 97 pp.
- Mehl, S., Aglen, A., Bogstad, B., Dingsør, G.E., Gjørseter, H., Godiksen, J., Johannesen, E., Korsbrekke, K., Staby, A., Wenneck, T. de Lange, Wienerroither, R., Murashko, P. A., and Russkikh, A. 2014. Fish investigations in the Barents Sea winter 2013-2014. IMR-PINRO Joint Report Series 2-2014, 73 pp.
- Mehl, S., and Yaragina, N. A. 1992. Methods and results in the joint PINRO-IMR stomach sampling program. In: Bogstad, B. and Tjelmeland, S. (eds.), *Interrelations between fish populations in the Barents Sea*. Proceedings of the fifth PINRO-IMR Symposium. Murmansk, 12–16 August 1991. Institute of Marine Research, Bergen, Norway, 5–16.
- Nilssen, K.T., Pedersen, O-P., Folkow, L. and Haug, T. 2000. Food consumption estimates of Barents Sea harp seals. NAMMCO Scientific Publications 2: 9-27.
- Prokhorova, T. (Ed.). 2013. Survey report from the joint Norwegian-Russian ecosystem survey in the Barents Sea and adjacent waters, August-October 2013. IMR/PINRO Joint Report Series, No. 4/2013. ISSN 1502-8828, 131 pp.
- Shevelev M.S., Mamylov V.S., Ratushny S.V., and E.N. Gavrilov, 1998. Technique of Russian bottom-trawl and acoustic surveys of the Barents Sea and how to improve them. NAFO Scientific Council Studies, No. 31, p.13-19.
- Trambachev, M.F. 1981. Young cod in the Barents Sea and Bear Island-Spitsbergen area in the autumn and winter 1978-1979. *Annls. biol., Copenh.*, 1981(1979), 36: 107-109.
- Yaragina, N. A. 2015. Short literature review on fishery-induced evolution in cod. WD 14, WKARCT 2015.
- Yaragina N.A., Kovalev Y. and Chetyrkin A. 2015. Hindcasting of NEA cod cannibalism for period 1946-1983. WD7, WKARCT 2015.
- Zaferman M.L. and Serebrov, L. I. 1984. On the instrumental methods for estimating bottom and demersal fish stocks in the Barents and Norwegian Seas. – In: *Reproduction and recruitment*

of Arctic cod. Reports of the 1st Soviet/Norwegian Symposium, Moscow, P. 359-370 (in Russian).

WD 1 WKBarFar 2021. Fall, J. Revision of NEA cod tuning indices from the Barents Sea Norwegian-Russian Winter Survey.

WD 2 WKBarFar 2021. Korsbrekke, K. Estimating variance for a trawl acoustic survey targeting spawning NEA cod.

WD6 WkBarFar 2021. Bogstad, B., and Nedreaas, K. NEA cod catch at age data revision.

WD 11 WKBarFar, 2021. Yaragina N.A., Kovalev Yu.A., Russkikh A.A. Scheme for estimating data on the Russian catches of Northeast Arctic cod.

WD 12 WKBarFar 2021. Kovalev Yu.A., Yaragina N.A. Northeast Arctic cod mean weight at age in stock. Problems and proposals.